

Product datasheet

Anti-Ago1 antibody ab5070

★★★★☆ 5 Abreviews 32 References 画像数 2

製品の概要

製品名	Anti-Ago1 antibody
製品の詳細	Rabbit polyclonal to Ago1
由来種	Rabbit
アプリケーション	適用あり: IP, IHC-P, WB
種交差性	交差種: Mouse, Drosophila melanogaster
免疫原	Synthetic peptide conjugated to KLH derived from within residues 1 - 100 of Fruit fly (Drosophila melanogaster) Ago1. Immunogenの所有権に関して (Peptide available as ab23538 .)
ポジティブ・コントロール	This antibody gave a positive signal in Drosophila embryo nuclear extract

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab5070** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
IP	★★★★☆	Use at an assay dependent concentration. PubMed: 18463631
IHC-P		Use at an assay dependent concentration. PubMed: 18528333

アプリケーション	Abreviews	特記事項
WB	★★★★☆	1/500 - 1/1000. Detects a band of approximately 100 kDa (predicted molecular weight: 110 kDa).

ターゲット情報

機能	Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs (miRNAs) or short interfering RNAs (siRNAs), and represses the translation of mRNAs which are complementary to them. Lacks endonuclease activity and does not appear to cleave target mRNAs. Also required for transcriptional gene silencing (TGS) of promoter regions which are complementary to bound short antisenRNA RNAs (agRNAs).
配列類似性	Belongs to the argonaute family. Ago subfamily. Contains 1 PAZ domain. Contains 1 Piwi domain.
細胞内局在	Cytoplasm > P-body.

画像



Western blot - Anti-Ago1 antibody (ab5070)

This image is courtesy of an anonymous Abreview

Anti-Ago1 antibody (ab5070) + *Drosophila* *Melanogaster* (Fruit Fly) Tissue Lysate at 15 μ g


Secondary

Sheep anti-rabbit HRP conjugated IgG
Polyclonal

Performed under reducing conditions.

Predicted band size: 110 kDa

Blot was blocked with 5% Milk for 1 hour at room temperature.



Anti-Ago1 antibody (ab5070) at 1 µg/ml +
Drosophila embryo nuclear extract (from
melanogaster embryos 0-12Hr) at 10 µg

Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-
Adsorbed (HRP) at 1/3000 dilution

Performed under reducing conditions.

Predicted band size: 110 kDa

Observed band size: 110 kDa

Additional bands at: 45 kDa. We are unsure
as to the identity of these extra bands.

Exposure time: 30 seconds

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors