

Product datasheet

Anti-CD31 antibody [TLD-3A12] ab64543

★★★★☆ 4 Abreviews 11 References 画像数 3

製品の概要

製品名	Anti-CD31 antibody [TLD-3A12]
製品の詳細	Mouse monoclonal [TLD-3A12] to CD31
由来種	Mouse
特異性	ab64543 recognises rat PECAM-1 (CD31). It also partially block the proliferative response of antigen-specific CD4+ T cells to antigen-presenting cells and relevant antigen.
アプリケーション	適用あり: WB, IHC-Fr, Flow Cyt, ICC/IF, ELISA 適用なし: IHC-P
種交差性	交差種: Rat 交差が予測される動物種: Pig, Rhesus monkey ▲ 非交差種: Human
免疫原	Activated, Lewis rat derived microglial cells
ポジティブ・コントロール	Rat spleen; activated microglial cells.
特記事項	This antibody has been shown to partially block the proliferative response of antigen-specific CD4+ T cells to antigen-presenting cells and relevant antigen.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	Preservative: 0.1% Sodium Azide Constituents: PBS, pH 7.4
精製度	Immunogen affinity purified
一次抗体 備考	This antibody has been shown to partially block the proliferative response of antigen-specific CD4+ T cells to antigen-presenting cells and relevant antigen.
ポリ/モノ	モノクローナル
クローン名	TLD-3A12
ミエローマ	Sp2
アイソタイプ	IgG1

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab64543** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	Use at an assay dependent concentration. Predicted molecular weight: 83 kDa.
IHC-Fr	★★★★★	1/10 - 1/100. A fixation with acetone or methanol is recommended. Do not fix with PFA.
Flow Cyt		Use 10µl for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration. PubMed: 20813915
ELISA		Use at an assay dependent concentration.

追加情報

Is unsuitable for IHC-P.

ターゲット情報

機能

Induces susceptibility to atherosclerosis (By similarity). Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions. Tyr-690 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes. Prevents phagocyte ingestion of closely apposed viable cells by transmitting 'detachment' signals, and changes function on apoptosis, promoting tethering of dying cells to phagocytes (the encounter of a viable cell with a phagocyte via the homophilic interaction of PECAM1 on both cell surfaces leads to the viable cell's active repulsion from the phagocyte. During apoptosis, the inside-out signaling of PECAM1 is somehow disabled so that the apoptotic cell does not actively reject the phagocyte anymore. The lack of this repulsion signal together with the interaction of the eat-me signals and their respective receptors causes the attachment of the apoptotic cell to the phagocyte, thus triggering the process of engulfment). Isoform Delta15 is unable to protect against apoptosis. Modulates BDKRB2 activation. Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in human umbilical cord vein cells (HUVEC).

組織特異性

Expressed on platelets and leukocytes and is primarily concentrated at the borders between endothelial cells. Isoform Long predominates in all tissues examined. Isoform Delta12 is detected only in trachea. Isoform Delta14-15 is only detected in lung. Isoform Delta14 is detected in all tissues examined with the strongest expression in heart. Isoform Delta15 is expressed in brain, testis, ovary, cell surface of platelets, human umbilical vein endothelial cells (HUVECs), Jurkat T-cell leukemia, human erythroleukemia (HEL) and U937 histiocytic lymphoma cell lines (at protein level).

配列類似性

Contains 6 Ig-like C2-type (immunoglobulin-like) domains.

ドメイン

The Ig-like C2-type domains 2 and 3 contribute to formation of the complex with BDKRB2 and in regulation of its activity.


翻訳後修飾

Phosphorylated on Ser and Tyr residues after cellular activation. In endothelial cells Fyn mediates mechanical-force (stretch or pull) induced tyrosine phosphorylation.

細胞内局在


Membrane. Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells and Cell junction. Localizes to the lateral border recycling compartment (LBRC) and recycles from the LBRC to the junction in resting endothelial cells.

画像



Immunohistochemical staining of rat brain cryosection with Mouse anti Rat CD31 (ab64543)


Immunohistochemistry (Frozen sections) - Anti-CD31 antibody [TLD-3A12] (ab64543)



ab64543 staining CD31 in rat brain tissue by Immunohistochemistry (Frozen sections). The sections were fixed in paraformaldehyde prior to blocking with 5% donkey serum, BSA 1%, Triton 1.5% for 2 hours at room temperature. The primary antibody was diluted 1/200 in PBS/5% Donkey serum and incubated with the sample for 12 hours at 4°C. An Alexa Fluor® 555-conjugated donkey anti-mouse polyclonal was used as the secondary antibody, diluted 1/200.

Immunohistochemistry (Frozen sections) - Anti-CD31 antibody [TLD-3A12] (ab64543)

This image is courtesy of Valley Lasry.



Immunohistochemical staining of rat spleen cryosection with Mouse anti Rat CD31 (ab64543)

Immunohistochemistry (Frozen sections) - Anti-CD31 antibody [TLD-3A12] (ab64543)

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors