

Product datasheet

Anti-CNTNAP3 antibody ab89720

2 References [画像数 2](#)

製品の概要

製品名	Anti-CNTNAP3 antibody
製品の詳細	Rabbit polyclonal to CNTNAP3
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, WB
種交差性	交差種: Human
免疫原	Synthetic peptide from a region within internal amino acids 612-661 SGPLGPFLVY CNMTADAAWT VVQHGGPDAV TLRGAPSGHP RSAVSFAYAA, of Human CNTNAP3 (NP_387504). Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	DU145 cell lysate This antibody gave a positive result when used in the following formaldehyde fixed cell lines: DU145

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: 0.09% Sodium azide Constituents: PBS, 2% Sucrose
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab89720** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 10 µg/ml.


アプリケーション	Abreviews	特記事項
----------	-----------	------

WB Use a concentration of 1 µg/ml. Predicted molecular weight: 141 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

ターゲット情報


配列類似性	Belongs to the neurexin family. Contains 2 EGF-like domains. Contains 1 F5/8 type C domain. Contains 1 fibrinogen C-terminal domain. Contains 4 laminin G-like domains.
細胞内局在	Secreted and Cell membrane.

画像



Immunocytochemistry/ Immunofluorescence - Anti-CNTNAP3 antibody (ab89720)

ICC/IF image of ab89720 stained DU145 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab89720 at 10µg/ml overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti- rabbit (ab96899) IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Anti-CNTNAP3 antibody (ab89720) at 1 µg/ml
(in 5% skim milk / PBS buffer) + DU145 cell
lysate at 10 µg

Secondary

HRP conjugated anti-Rabbit IgG at 1/50000
dilution

Predicted band size: 141 kDa

Observed band size: ~144 kDa

Additional bands at: 90 kDa. We are unsure
as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors