


機能	Mediates the formation of prostaglandins from arachidonate. May have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity.
パスウェイ	Lipid metabolism; prostaglandin biosynthesis.
配列類似性	Belongs to the prostaglandin G/H synthase family. Contains 1 EGF-like domain.
翻訳後修飾	S-nitrosylation by NOS2 (iNOS) activates enzyme activity. S-nitrosylation may take place on different Cys residues in addition to Cys-561.
細胞内局在	Microsome membrane. Endoplasmic reticulum membrane.

Anti-COX2 / Cyclooxygenase 2 antibody [SP21] 画像



ab16701 at 1/100 dilution staining COX2 / Cyclooxygenase 2 in human breast carcinoma by Immunohistochemistry (FFPE-sections).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - COX2 / Cyclooxygenase 2 antibody [SP21] (ab16701)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors