

Product datasheet

Anti-Calponin antibody [CALP] ab700

★★★★★ 3 Abreviews 14 References 画像数 2

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-Calponin antibody [CALP] |
| 製品の詳細 | Mouse monoclonal [CALP] to Calponin |
| アプリケーション | 適用あり: IHC-Fr, Flow Cyt, ICC/IF, IHC-FoFr, IHC-P |
| 種交差性 | 交差種: Rat, Human |
| 免疫原 | Crude human uterus extract |
| ポジティブ・コントロール | Myoepithelium in normal breast glands. |

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | Preservative: Proprietary, not sodium azide or thimerosal. Constituents: PBS, Tissue Culture Supernatant. pH 7.3, Protein carrier. |
| ポリ/モノ | モノクローナル |
| クローン名 | CALP |
| ミエローマ | unknown |
| アイソタイプ | IgG1 |
| 軽鎖の種類 | kappa |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab700** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|------|
| | | |

| アプリ ケー ション | Abreviews | 特記事項 |
|------------------|-----------|---|
| IHC- Fr | | 1/100 - 1/400. ABC system. 30mins at RT. Calponin has been found to be useful as a marker for myoepithelial and basal lamina in differentiating microinvasive from in situ ductal carcinomas of the breast. Calponin has also been useful in malignant myoepithelium and pleomorphic adenoma of salivary gland as well as a useful marker for fine needle aspirates of papillary breast lesions. |
| Flow Cyt | | Use at an assay dependent concentration. PubMed: 20618694 |
| ICC/IF | ★★★★★ | Use at an assay dependent concentration. |
| IHC- FoFr | | Use at an assay dependent concentration. PubMed: 20138661 |
| IHC-P | ★★★★★ | 1/100 - 1/400. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. ABC system. 30mins at RT. |

ターゲット情報

| | |
|-------|--|
| 機能 | Thin filament-associated protein that is implicated in the regulation and modulation of smooth muscle contraction. It is capable of binding to actin, calmodulin, troponin C and tropomyosin. The interaction of calponin with actin inhibits the actomyosin Mg-ATPase activity. |
| 組織特異性 | Smooth muscle, and tissues containing significant amounts of smooth muscle. |
| 配列類似性 | Belongs to the calponin family. Contains 3 calponin-like repeats. Contains 1 CH (calponin-homology) domain. |


画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Calponin antibody [CALP] (ab700)

This image is courtesy of an Abreview submitted by Dr Mal Niladri

ab700 at a 1/50 dilution staining rat heart tissue sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). The tissue was formaldehyde fixed prior to incubation with the antibody overnight at 4°C. An Alex-Fluor® 488 conjugated goat anti-mouse antibody was used as the secondary. The image was taken with a confocal laser scanning microscope with an additional Differential Interference Contrast (DIC) mode and shows stained Calponin (smooth muscle cells-green) and counterstained nuclei (DAPI-blue) (left panel). DIC (phase) image of the same tissue (right panel) confirm the localization of the calponin staining in the arterial smooth muscle in the heart tissue with very little background.



Immunocytochemistry/ Immunofluorescence - Calponin antibody [CALP] (ab700)

This image is a courtesy of Anonymous Abreview

ab700 staining Calponin in rat Smooth muscle cells from mesenteric artery by Immunocytochemistry/ Immunofluorescence. Cells were fixed with 4% paraformaldehyde in physiological saline solution (PSS) for 4 min at 4°C and permeabilized with 0.3% Triton x100 before blocking with 2% BSA was done for 30 minutes at 20°C. Samples were incubated with primary antibody (1/300: in PSS with 2%BSA and 0.3% Triton X-100) for 14 hours at 4°C. An Alexa Fluor®488-conjugated chicken polyclonal to mouse IgG was used as secondary antibody at 1/400 dilution. Image shows combined fluorescent and transmitted light images. Hoechst 33342 was used to stain the cell nuclei (blue).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors