

Product datasheet

Anti-Cardiac Troponin T antibody [1C11] ab8295

★★★★★ 6 Abreviews 53 References 画像数 5

製品の概要

製品名	Anti-Cardiac Troponin T antibody [1C11]
製品の詳細	Mouse monoclonal [1C11] to Cardiac Troponin T
由来種	Mouse
アプリケーション	適用あり: Flow Cyt, IHC-Fr, WB, ELISA, IHC-FoFr, IP, IHC-P, Sandwich ELISA, ICC/IF
種交差性	交差種: Mouse, Rat, Dog, Human
免疫原	Other Immunogen Type corresponding to Human Cardiac Troponin T aa 171-190. Database link: P45379
特記事項	This antibody detects Troponin T in human cardiac muscle. No cross-reaction with skeletal troponin T, cTnl and TnC. This product was changed from ascites to tissue culture supernatant on 17th October 2017 and product received after this date will be from tissue culture supernatant.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	PBS with 0.1% sodium azide, pH 7.4
精製度	Tissue culture supernatant
一次抗体 備考	This antibody detects Troponin T in human cardiac muscle. No cross-reaction with skeletal troponin T, cTnl and TnC.
ポリ/モノ	モノクローナル
クローン名	1C11
ミエローマ	Sp2/0
アイソタイプ	IgG1
軽鎖の種類	unknown

アプリケーション


Our [Abpromise guarantee](#) covers the use of **ab8295** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
Flow Cyt	★★★★★	Use at an assay dependent concentration. PubMed: 18593559
IHC-Fr	★★★★★	Use at an assay dependent concentration. PubMed: 17916803
AP		Use at an assay dependent concentration.
WB	★★★★☆	Use at an assay dependent concentration. Predicted molecular weight: 35.4 kDa. PubMed: 17916803
ELISA		Use at an assay dependent concentration.
IHC-FoFr		Use at an assay dependent concentration. PubMed: 19859557
IP	★★★★★	Use at an assay dependent concentration.
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
Sandwich ELISA		Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Mouse monoclonal [9G6] to Cardiac Troponin T (HRP) (ab8296) . Can be used as Capture or Detection antibody.
ICC/IF	★★★★★	Use at an assay dependent concentration. PubMed: 18593559

ターゲット情報


機能	Troponin T is the tropomyosin-binding subunit of troponin, the thin filament regulatory complex which confers calcium-sensitivity to striated muscle actomyosin ATPase activity.
組織特異性	Heart. The fetal heart shows a greater expression in the atrium than in the ventricle, while the adult heart shows a greater expression in the ventricle than in the atrium. Isoform 6 predominates in normal adult heart. Isoforms 1, 7 and 8 are expressed in fetal heart. Isoform 7 is also expressed in failing adult heart.
関連疾患	Defects in TNNT2 are the cause of cardiomyopathy familial hypertrophic type 2 (CMH2) [MIM:115195]. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death. Defects in TNNT2 are the cause of cardiomyopathy dilated type 1D (CMD1D) [MIM:601494]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death. Defects in TNNT2 are the cause of cardiomyopathy familial restrictive type 3 (RCM3) [MIM:612422]. Restrictive cardiomyopathy is a heart disorder characterized by impaired filling of the ventricles with reduced diastolic volume, in the presence of normal or near normal wall thickness and systolic function.
配列類似性	Belongs to the troponin T family.



Immunohistochemistry (Frozen sections) - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

This image is courtesy of an anonymous Abreview.


ab8295 staining Cardiac Troponin T in mouse cardiac tissue sections by Immunohistochemistry (IHC-Fr - frozen sections). Tissue was fixed with formaldehyde, permeabilized with 0.5% Tx100 in PBS and blocked with 1% BSA for 90 minutes at 20°C. Samples were incubated with primary antibody (1/200 in 1% goat serum, 10% BSA + 0.1% Tx100 PBS) for 16 hours at 4°C. An Alexa Fluor® 488-conjugated goat anti-mouse IgG polyclonal (1/400) was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

This image is courtesy of an anonymous Abreview.

ab8295 staining Cardiac Troponin T in human pluripotent stem cell derived cardiomyocytes by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized with saponin and blocked with 5% serum for 15 minutes at room temperature. Samples were incubated with primary antibody (1/800) for 1 hour at 21°C. An Alexa Fluor®-568 goat anti-mouse IgG monoclonal (1/1000) was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Cardiac Troponin T antibody [1C11] (ab8295)


This image is courtesy of an anonymous Abreview.

ab8295 staining Cardiac Troponin T in the human iPS cell derived from cardiomyocytes by ICC/IF

(Immunocytochemistry/immunofluorescence).

Cells were fixed with paraformaldehyde, permeabilized with 0.2% triton X in PBS and blocked with 1% BSA for 18 hours at 4°C.

Samples were incubated with primary antibody (1/100 in 1% BSA/PBS) for 18 hours at 4°C. An Alexa Fluor® 488-conjugated goat anti-mouse IgG polyclonal (1/500) was used as the secondary antibody.



Sandwich ELISA - Anti-Cardiac Troponin T antibody [1C11] (ab8295)


Calibration curves for sandwich cTnT

fluoroimmunoassay with different animal TnTs as antigen. (dark blue) canine, (blue/grey)

human, (grey) mouse, (black) rat. Monoclonal antibodies: capture, ab8295 [clone 1C11], 1

µg/well, detection ab1454 [clone 7E7], 200 ng/well. Assay time, 30 min at room

temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

Ab8295 staining human normal heart. Staining is localised to the cytoplasm.

Left panel: with primary antibody at 1 ug/ml.

Right panel: isotype control.

Sections were stained using an automated system DAKO Autostainer Plus , at room temperature. Sections were rehydrated and antigen retrieved with the Dako 3-in-1 AR buffer citrate pH 6.0 in a DAKO PT Link.

Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS), then incubated with primary antibody for 20 minutes, and detected with Dako Envision Flex amplification kit for 30 minutes.

Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors