


Product datasheet

Anti-EGR4 antibody ab50636

画像数 1

製品の概要

製品名	Anti-EGR4 antibody
製品の詳細	Rabbit polyclonal to EGR4
由来種	Rabbit
アプリケーション	適用あり: ELISA, WB
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat, Cow, Dog, Drosophila melanogaster, Zebrafish 
免疫原	Synthetic peptide (Human)
ポジティブ・コントロール	Jurkat cell lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None Constituents: 2% Sucrose, PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab50636** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration. Peptide ELISA titre: 1/312500.

アプリ ケーショ ン	Abreviews	特記事項
------------------	-----------	------


WB Use a concentration of 0.0625 µg/ml. Detects a band of approximately 46 kDa (predicted molecular weight: 51 kDa). Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

## ターゲット情報

**関連性** EGR4(Early Growth Response protein 4) is a zinc-finger transcription factor that activates the transcription of genes whose products are required for mitogenesis and differentiation. EGR4 is closely related to nerve growth factor-induced clone A (NGF1A or EGR1), EGR2, and EGR3. These four early response (immediate early) proteins all contain very similar zinc-finger DNA binding domains and five highly homologous subdomains. EGR4 functionally cooperate with NFAT proteins and induce expression of IL2 and TNFalpha. Early growth response proteins (EGR) and nuclear factors of activated T cells (NFAT) form heterodimers and regulate proinflammatory cytokine gene expression.

**細胞内局在** Nuclear

## 画像



Anti-EGR4 antibody (ab50636) at 0.0625 µg/ml + Jurkat cell lysate at 10 µg

### Secondary

HRP conjugated anti-Rabbit IgG at 1/50,000 - 1/100,000 dilution.

**Predicted band size:** 51 kDa

**Observed band size:** 46 kDa

The primary antibody was diluted in 5% skim milk/PBS buffer.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors