

Product datasheet

Anti-FGF17 antibody [MM0284-5H27] ab89270

画像数 1

製品の概要

| | |
|----------|-----------------------------------------|
| 製品名 | Anti-FGF17 antibody [MM0284-5H27] |
| 製品の詳細 | Mouse monoclonal [MM0284-5H27] to FGF17 |
| 由来種 | Mouse |
| アプリケーション | 適用あり: Sandwich ELISA, WB, IHC-Fr |
| 種交差性 | 交差種: Human |
| 免疫原 | Recombinant Human FGF17. |

製品の特性

| | |
|--------|---------------------------------------------------------------------------------------------------------|
| 製品の状態 | Lyophilised:Reconstitute the antibody with 200 µl sterile PBS. |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles. |
| バッファー | Preservative: None Constituents: PBS |
| 精製度 | Protein G purified |
| ポリ/モノ | モノクローナル |
| クローン名 | MM0284-5H27 |
| アイソタイプ | IgG2 |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab89270** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


| アプリケーション | Abreviews | 特記事項 |
|----------------|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sandwich ELISA | | Use a concentration of 1 µg/ml. Can be paired for Sandwich ELISA with Rabbit polyclonal to FGF17 (ab9990) . For sandwich ELISA, use this antibody as Capture at 1µg/ml with ab9990 as Detection. |
| WB | | 1/500 - 1/1000. Predicted molecular weight: 25 kDa. |

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---------------|
| IHC-Fr | | 1/50 - 1/200. |

ターゲット情報

| | |
|-------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 機能 | Plays an important role in the regulation of embryonic development and as signaling molecule in the induction and patterning of the embryonic brain. Required for normal brain development. |
| 組織特異性 | Preferentially expressed in the embryonic brain. |
| 配列類似性 | Belongs to the heparin-binding growth factors family. |
| 発生段階 | Detected in embryos at E14.5, but not at E10.5 and E19.5. Preferentially expressed in the neuroepithelia of the isthmus and septum of the embryonic brain at E14.5. |
| 細胞内局在 | Secreted. |

画像



Standard Curve for FGF17 (Analyte: [ab9991](#)) dilution range 1pg/ml to 1ug/ml using Capture Antibody [Mouse monoclonal \[MM0284-5H27\]](#) to FGF17 ([ab89270](#)) at 1ug/ml and Detector Antibody [Rabbit polyclonal to FGF17 \(ab9990\)](#) at 0.5ug/ml

Sandwich ELISA - FGF17 antibody [[MM0284-5H27](#)]
([ab89270](#))

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors