


Product datasheet

Anti-FOXK1 antibody ab85999

画像数 3

製品の概要

製品名	Anti-FOXK1 antibody
製品の詳細	Rabbit polyclonal to FOXK1
由来種	Rabbit
アプリケーション	適用あり: WB, IP, IHC-P
種交差性	交差種: Mouse, Human 交差が予測される動物種: Horse, Guinea pig, Cow, Pig, Chimpanzee, Rhesus monkey, Gorilla 
免疫原	Synthetic peptide corresponding to a region between residues 683 and 733 of human FOXK1 using the numbering given in entry NP_001032242.1.
ポジティブ・コントロール	Whole cell lysate from HeLa cells

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.09% Sodium Azide Constituents: 0.1% BSA, Tris buffered saline
精製度	Immunogen affinity purified
特記事項(精製)	ab85999 was affinity purified using an epitope specific to FOXK1 immobilized on solid support.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab85999** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/2000 - 1/10000. Predicted molecular weight: 75 kDa.
IP		Use at 2-5 µg/mg of lysate.
IHC-P		1/200 - 1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## ターゲット情報


機能	Transcriptional regulator that binds to the upstream enhancer region (CCAC box) of myoglobin gene. Has a role in myogenic differentiation and in remodeling processes of adult muscles that occur in response to physiological stimuli.
組織特異性	Expressed both developing and adult tissues. In adults, significant expression is seen in tumors of the brain, colon and lymph node.
配列類似性	Contains 1 FHA domain. Contains 1 fork-head DNA-binding domain.
細胞内局在	Nucleus.

## 画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma (left) and mouse teratoma (right) tissues labelling FOXX1 with ab85999 at 1/1000 (0.2µg/ml) and 1/200 (1µg/ml). Detection: DAB.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXX1 antibody (ab85999)



Western blot - FOXX1 antibody (ab85999)

**All lanes :** Anti-FOXX1 antibody (ab85999) at 0.04 µg/ml

**Lane 1 :** HeLa whole cell lysate at 50 µg

**Lane 2 :** HeLa whole cell lysate at 15 µg


**Lane 3 :** HeLa whole cell lysate at 5 µg

**Lane 4 :** 293T whole cell lysate at 50 µg

Developed using the ECL technique.

**Predicted band size:** 75 kDa

**Exposure time:** 30 seconds



Immunoprecipitation - FOXX1 antibody (ab85999)

Detection of FOXX1 by Western blot of Immunoprecipitate.

ab85999, at 1 µg/ml, staining FOXX1 in HeLa whole cell lysate immunoprecipitated using ab85999 at 3 µg/mg lysate (1 mg/IP; 20% of IP loaded/lane).

Detection: Chemiluminescence with an exposure time of 3 seconds.

**All lanes :** Anti-FOXX1 antibody (ab85999) at 1 µg/ml

**Lane 1 :** HeLa whole cell lysate

**Lane 2 :** Control IgG

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors