


Product datasheet

Anti-GARNL3 antibody ab122015

画像数 3

製品の概要

製品名	Anti-GARNL3 antibody
製品の詳細	Rabbit polyclonal to GARNL3
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Mouse 
免疫原	amino acids 737-828 (LVQPSASDFQ FCWNQAPYAI VCAFPYLLAF TTDSMEIRLV VNGNLVHTAV VPQLQLVASR SDIYFTATAA VNEVSSGGSS KGASARNSPQ TP) of Human GARNL3 (Q5VWW2). Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	Human nasopharynx tissue.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol
精製度	Immunogen affinity purified
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab122015** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.
ICC/IF		Use a concentration of 1 - 4 µg/ml. Recommend PFA Fixation and Triton X-100 treatment
IHC-P		1/20 - 1/50. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.

ターゲット情報


配列類似性	<p>Belongs to the GARNL3 family.</p> <p>Contains 1 CNH domain.</p> <p>Contains 1 Rap-GAP domain.</p>
-------	--

画像




Immunofluorescent staining of Human cell line U-251MG shows positivity in vesicles. Recommended concentration of ab122015 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-GARNL3 antibody (ab122015)



ab122015 staining GARNL3 in paraffin-embedded Human nasopharynx tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GARNL3 antibody (ab122015)



Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10

Lane 2: Negative control (vector only transfected HEK293T lysate)

Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells.

Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10

Lane 2: Negative control (vector only transfected HEK293T lysate)

Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells.

Western blot - Anti-GARNL3 antibody (ab122015)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors