

Product datasheet

Anti-GATA5 antibody ab48820

★★★★☆ 1 Abreviews 1 References 画像数 2

製品の概要

製品名	Anti-GATA5 antibody
製品の詳細	Rabbit polyclonal to GATA5
特異性	This antibody recognizes GATA5
アプリケーション	適用あり: ELISA, WB, ICC/IF
種交差性	交差種: Mouse 交差が予測される動物種: Rat, Human
免疫原	A region within synthetic peptide: LNSESSATTL KAESSLASPV CAGPTITSQA SSPADESLAS SHLEFKFEPE, corresponding to C terminal amino acids 325-374 of Human GATA5 Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	SP2/0 lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: None Constituents: 2% Sucrose, PBS
精製度	Protein A purified
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab48820** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
ELISA		1/1.5625e+006.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1.25 µg/ml. Predicted molecular weight: 44 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.
ICC/IF	★★★★☆	1/100.

ターゲット情報

機能	Binds to the functionally important CEF-1 nuclear protein binding site in the cardiac-specific slow/cardiac troponin C transcriptional enhancer. May play an important role in the transcriptional program(s) that underlies smooth muscle cell diversity.
配列類似性	Contains 2 GATA-type zinc fingers.
細胞内局在	Nucleus.

画像



Anti-GATA5 antibody (ab48820) at 1.25 µg/ml
+ SP2/0 lysate at 10 µg with 5% skimmed milk
in PBS.


Secondary

HRP conjugated anti-Rabbit IgG at 1/50000
dilution

Predicted band size : 44 kDa

Observed band size : 41 kDa

Gel concentration 12%



Immunocytochemistry/ Immunofluorescence - Anti-GATA5 antibody (ab48820)

Image courtesy of an anonymous Abreview.

ab48820 staining GATA5 in murine epicardial explant culture by Immunocytochemistry/ Immunofluorescence.

Cells were fixed in formaldehyde, permeabilized using 0.5% Triton X-100 in PBS, blocked with 1% goat serum 10% BSA 0.1% Triton X-100 PBS and then incubated with ab48820 at a 1/100 dilution for 16 hours at 4°C. The secondary used was an Alexa-Fluor 594 conjugated goat anti-rabbit IgG (H+L) used at a 1/200 dilution.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors