

Product datasheet

Anti-Hepatitis B Virus Surface Antigen (Ad/Ay) antibody (Biotin) ab20778

製品の概要

| | |
|----------|---|
| 製品名 | Anti-Hepatitis B Virus Surface Antigen (Ad/Ay) antibody (Biotin) |
| 製品の詳細 | Goat polyclonal to Hepatitis B Virus Surface Antigen (Ad/Ay) (Biotin) |
| 標識 | Biotin |
| 特異性 | Non-reactive with normal human serum. Monospecific, reacts only with Hepatitis B virus. |
| アプリケーション | 適用あり: ICC/IF, ELISA |
| 免疫原 | Mixture of subtypes ad & ay. Hepatitis B surface antigen purified from human serum |
| 特記事項 | Covalently coupled with the N-Hydroxysuccinimide ester of biotin under mild conditions to give a high degree of substitution. |

製品の特性

| | |
|----------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | Preservative: 0.1% Sodium Azide Constituents: 10mM PBS, pH 7.2 |
| 精製度 | IgG fraction |
| 特記事項(精製) | Purified IgG fraction of the antiserum covalently coupled with the N-Hydroxysuccinimide ester of Biotin. |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab20778** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|------|
| ICC/IF | | |

ELISA

追加情報

ELISA: 1/1:1000.

IF: Use at an assay dependent dilution. Use Avidin and streptavidin amplifications systems.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性

Hepatitis B Virus (HBV) infection induces a disease state characterised by liver damage, inflammation and viral persistence. Infection also increases the risk of hepatocellular carcinoma. HBV belongs to the Hepadnaviridae family of viruses. Its genome consists of partially double stranded circular DNA. The DNA is enclosed in a nucleocapsid, or core antigen (HBcAg), which is surrounded by a spherical envelope (surface antigen or HBsAg). The core antigen shares its sequences with the e antigen (HBeAg) but no cross reactivity between the two proteins has been observed. The HBV genome also encodes a DNA polymerase that also acts as a reverse transcriptase. Hepatitis B infection is normally diagnosed from serological tests that detect HBsAg but as the disease progresses this antigen may no longer be present in the blood and tests for HBcAg are used. If HBsAg can be detected in the blood for longer than six months, chronic hepatitis B is diagnosed. The antigenic determinant of the protein moiety of the HBsAg determines specific characteristics of different serotypes and provides the basis of immunodetection. HBsAg has antigenic heterogeneity, specifically, two pairs of sub specific determinants, d/y and w/r allow the following combinations: adw, ayw, adr, ayr.

細胞内局在

Virion membrane

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors