

## Product datasheet

# Human NADPH oxidase 4 peptide ab104391

### 製品の概要

---

製品名 Human NADPH oxidase 4 peptide

### 製品の詳細

---

由来 Synthetic

### アミノ酸配列

生物種 Human

### 特性

---

Our [Abpromise guarantee](#) covers the use of **ab104391** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

精製度 70 - 90% by HPLC.

製品の状態 Liquid

### 備考

- First try to dissolve a small amount of peptide in either water or buffer. The more charged residues on a peptide, the more soluble it is in aqueous solutions.
- If the peptide doesn't dissolve try an organic solvent e.g. DMSO, then dilute using water or buffer.
- Consider that any solvent used must be compatible with your assay. If a peptide does not dissolve and you need to recover it, lyophilise to remove the solvent.
- Gentle warming and sonication can effectively aid peptide solubilisation. If the solution is cloudy or has gelled the peptide may be in suspension rather than solubilised.
- Peptides containing cysteine are easily oxidised, so should be prepared in solution just prior to use.

### 前処理および保存

---

保存方法および安定性 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Information available upon request.

## 関連情報

---

<b>機能</b>	Constitutive NADPH oxidase which generates superoxide intracellularly upon formation of a complex with CYBA/p22phox. Regulates signaling cascades probably through phosphatases inhibition. May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity. May regulate insulin signaling cascade. May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFkB. May produce superoxide in the nucleus and play a role in regulating gene expression upon cell stimulation. Isoform 3 is not functional. Isoform 4 displays an increased activity. Isoform 5 and isoform 6 display reduced activity.
<b>組織特異性</b>	Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells.
<b>配列類似性</b>	Contains 1 FAD-binding FR-type domain. Contains 1 ferric oxidoreductase domain.
<b>発生段階</b>	Expressed in fetal kidney and fetal liver.
<b>翻訳後修飾</b>	Isoform 3 and isoform 4 are N-glycosylated. Isoform 4 glycosylation is required for its proper function.
<b>細胞内局在</b>	Endoplasmic reticulum membrane. Cell membrane. Cell junction > focal adhesion. Nucleus. May localize to plasma membrane and focal adhesions. According to PubMed:15927447, may also localize to the nucleus.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors