abcam

Product datasheet

Human TNF alpha ELISPOT Kit (with un-coated plates) ab46566

製品の概要

製品名	Human TNF alpha ELISPOT Kit (with un-coated plates)
検出方法	Colorimetric
サンプルの種類	Suspension cells
アッセイタイプ	Sandwich (qualitative)
ステップ	Multiple steps standard assay
種交差性	**交差種:** Human, Macaque monkey
製品の概要	The Human TNF alpha ELISPOT Kit (with un-coated plates) is designed to enumerate cytokine producing cells in a single cell suspension. This method has the advantage of requiring a minimum of in-vitro manipulations allowing cytokine production analysis as close as possible to in-vivo conditions in a highly specific way. This technique is designed to determine the frequency of cytokine producing cells under a given stimulation, and the follow-up of such frequency during a treatment and/or a pathological state. Elispot assay constitutes an ideal tool in the TH1 / TH2 response, vaccine development, viral infection monitoring and treatment, cancerology, infectious diseases, autoimmune diseases and transplantation. This ELISPOT assay is based on sandwich immuno-enzyme technology. Cell secreted cytokines or soluble molecules are captured by coated antibodies avoiding diffusion in supernatant, protease degradation or binding on soluble membrane receptors. After cell removal, the captured cytokines are revealed by tracer antibodies and appropriate conjugates. **Principal:** After cell stimulation, locally produced cytokines are captured by a specific monoclonal antibody. After cell lysis, trapped cytokine molecules are revealed by a secondary biotinylated detection antibody, which is in turn recognised by streptavidin conjugated to alkaline phosphatase. PVDF-bottomed-well plates are then incubated with BCIP/NBT substrate. Colored "purple" spots indicate cytokine production by individual cells. Recognizes natural human TNF alpha
アプリケーション	**適用あり:** ELISpot
試験プラットフォーム	Microplate

製品の特性

保存方法 Store at +4°C. Please refer to protocols.

内容	5 x 96 tests	10 x 96 tests
96 PVDF-bottomed-well plates.	4 units	10 units
Bovine Serum Albumin	1 x 1g	2 x 1g
TNFα Capture antibody (clone B-F7)	1 x 500µl	2 x 500µl
Ready-to-use BCIP/NBT substrate buffer	1 x 50ml	2 x 50ml
Streptavidin - Alkaline Phosphatase conjugated	1 x 50µl	2 x 50µl
TNFα Biotinylated detection antibody (clone B-C7)	1 vial	2 vials

機能 Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation.

関連疾患 Genetic variations in TNF are a cause of susceptibility psoriatic arthritis (PSORAS) [MIM:607507]. PSORAS is an inflammatory, seronegative arthritis associated with psoriasis. It is a heterogeneous disorder ranging from a mild, non-destructive disease to a severe, progressive, erosive arthropathy. Five types of psoriatic arthritis have been defined: asymmetrical oligoarthritis characterized by primary involvement of the small joints of the fingers or toes; asymmetrical arthritis which involves the joints of the extremities; symmetrical polyarthritis characterized by a rheumatoidlike pattern that can involve hands, wrists, ankles, and feet; arthritis mutilans, which is a rare but deforming and destructive condition; arthritis of the sacroiliac joints and spine (psoriatic spondylitis).

配列類似性 Belongs to the tumor necrosis factor family.

翻訳後修飾 The soluble form derives from the membrane form by proteolytic processing.
The membrane form, but not the soluble form, is phosphorylated on serine residues. Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1.
O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid.

細胞内局在 Secreted and Cell membrane.

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab46566の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISpot		Use at an assay dependent dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit **https://www.abcam.co.jp/abpromise** or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors