

Product datasheet

Anti-IGF2BP1 antibody ab107205

1 Abreviews 2 References 画像数 1

製品の概要

製品名	Anti-IGF2BP1 antibody
製品の詳細	Rabbit polyclonal to IGF2BP1
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Synthetic peptide conjugated to KLH, corresponding to a region within C terminal amino acids 515-542 of Human IGF2BP1 (NP_006537.3).
ポジティブ・コントロール	K562 cell line lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at 4°C (up to 6 months). Store at -20°C long term.
バッファー	Preservative: 0.09% Sodium Azide Constituents: PBS
精製度	Immunogen affinity purified
特記事項(精製)	ab107205 is purified through a protein A column, followed by peptide affinity purification.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab107205** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/100 - 1/500. Predicted molecular weight: 63 kDa.

ターゲット情報

機能

RNA-binding factor that affects mRNA nuclear export, localization, stability and translation. Component of the CRD-mediated complex that promotes MYC mRNA stabilization. Regulates mRNA stability during the integrated cellular stress response (ISR) in stress granules (SGs). Stabilizes the BTRC/FBW1A mRNA from degradation by disrupting miRNA-dependent interaction with AGO2. Identified in a HCV IRES-mediated translation complex, that enhances translation at the Hepatitis C virus (HCV) RNA-replicon via the internal ribosome entry site (IRES), but does not affect 5'cap-dependent translation. Acts as a HIV-1 retrovirus restriction factor that reduces HIV-1 assembly by inhibiting viral RNA packaging, assembly and processing of HIV-1 GAG protein on cellular membranes. Binds to mRNAs in stress granules (SGs). Binds to the stem-loop IV of the 5'-UTR and to the variable region and the poly(U-C) motif of the 3'-UTR of the HCV RNA-replicon. Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNA and regulates its subcellular localization and translation. Binds both to the coding region mRNA stability determinant (CRD) and to AU-rich sequences in the 3'-UTR of the MYC and CD44 mRNAs and stabilizes these mRNAs. Binds to the fourth and fifth exons of the oncofetal H19 and neuron-specific TAU mRNAs and regulates their localizations. Binds to the adenine-rich autoregulatory sequence (ARS) 5'-UTR of the PABPC1 mRNA and is involved in its translational repression. The RNA-binding activity to ARS is stimulated by PABPC1. Binds to the coding sequence region of BTRC/FBW1A mRNA and mediates stabilization of BTRC/FBW1A and MYC mRNAs in response to beta-catenin signaling. Binding to RNA employs a cooperative, sequential mechanism of homo- or heterodimerisation. Also involved in growth or survival of lung-cancer cells. Protects the MYC and MDR-1 mRNAs from cleavage by a endoribonuclease, thus prolonging their stabilities (By similarity). Binds to the 3'-UTR axonal localization signal (ALS) of TAU mRNA (By similarity). Binds to a conserved 54-nucleotide element in the 3'-UTR of the beta actin mRNA known as the 'zipcode' (By similarity). Promotes translocation of the beta-actin mRNA to dendrites (By similarity). May act as a regulator of mRNA transport to activated synapses in response to synaptic activity.

組織特異性

Expressed in fetal liver, fetal lung, fetal kidney, fetal thymus, fetal placenta, fetal follicles of ovary, gonocytes of testis, oocytes, spermatogonia and semen (at protein level). Expressed in testicular and lung cancer (at protein level). Expressed in kidney, prostate, trachea, testis and lung cancer.

配列類似性

Belongs to the RRM IMP/VICKZ family.
Contains 4 KH domains.
Contains 2 RRM (RNA recognition motif) domains.

ドメイン

The third and fourth KH domains encompass the protein dimerization motif and are necessary and sufficient for RNA binding. The four KH domains are important for granule formation and SGs targeting. Contains two nuclear export signals, situated within the second and fourth KH domains. The four KH domains are important to suppress HIV-1 infectivity.

翻訳後修飾


Phosphorylated. Phosphorylation may influence mRNA translation.

細胞内局在

Nucleus. Cytoplasm. Cell projection > lamellipodium. Cell projection > dendrite. Cell projection > dendritic spine. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Targeted to stress granules (SGs), but not processing bodies (PBs), during cellular stress. Colocalizes with G3BP1 and TIAL1 in SGs. Colocalizes with HIV-1 GAG at the cell edges. Found in lamellipodia of the leading edge, in the perinuclear region, and beneath the plasma membrane. The subcytoplasmic localization is cell specific and regulated by cell contact and growth. Colocalized with H19 RNA at lamellipodia. Colocalized with CD44 mRNA in RNP granules. Nuclear export is mediated by XPO1/CRM1. In motile cells, is transported towards the leading edge into the cortical region of the lamellipodia where it is connected to microfilaments (By similarity). Present in the form of granules and into F-actin-rich protrusion of dendrites, spines and subsynaptic sites (By similarity). Colocalizes with beta-actin mRNA in dendrites and

spines (By similarity). Exhibited rapid, bidirectional movements in dendrites and spines (By similarity). Neuronal depolarization by KCl induces its rapid efflux from the cell body into dendrites.

画像



Anti-IGF2BP1 antibody (ab107205) at 1/100 dilution + K562 cell line lysate at 35 µg

Predicted band size: 63 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors