

## Product datasheet

# Anti-Interferon gamma antibody [DB-1] ab22543

### 製品の概要

製品名	Anti-Interferon gamma antibody [DB-1]
製品の詳細	Mouse monoclonal [DB-1] to Interferon gamma
特異性	This antibody reacts with both mouse and rat Interferon gamma. Clone DB-1 exhibits high affinity binding to the natural and recombinant forms and effectively neutralises the anti viral activity of IFN-gamma. It also blocks the induction of La antigen expression on murine keratinocytes.
アプリケーション	<b>適用あり:</b> ELISA, Blocking, Neutralising
種交差性	<b>交差種:</b> Mouse, Rat
免疫原	Recombinant full length rat IFN gamma

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.09% Sodium Azide Constituents: PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	DB-1
アイソタイプ	IgG1

### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab22543** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		
Blocking		

Neutralising

#### 追加情報

BL: Use at an assay dependent dilution  
ELISA: Use at a concentration of 10 µg/ml, 100ul/well as coating antibody  
Neut: Use at an assay dependent dilution.

Not tested in other applications.  
Optimal dilutions/concentrations should be determined by the end user.

#### ターゲット情報

##### 機能

Produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.

##### 組織特異性

Released primarily from activated T lymphocytes.

##### 関連疾患

In Caucasians, genetic variation in IFNG is associated with the risk of aplastic anemia (AA) [MIM:609135]. AA is a rare disease in which the reduction of the circulating blood cells results from damage to the stem cell pool in bone marrow. In most patients, the stem cell lesion is caused by an autoimmune attack. T-lymphocytes, activated by an endogenous or exogenous, and most often unknown antigenic stimulus, secrete cytokines, including IFN-gamma, which would in turn be able to suppress hematopoiesis.

##### 配列類似性

Belongs to the type II (or gamma) interferon family.

##### 翻訳後修飾

Proteolytic processing produces C-terminal heterogeneity, with proteins ending alternatively at Gly-150, Met-157 or Gly-161.

##### 細胞内局在

Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors