


Product datasheet

# Anti-Ionotropic Glutamate receptor 4 antibody ab77407

1 References [画像数 1](#)

製品の概要

製品名	Anti-Ionotropic Glutamate receptor 4 antibody
製品の詳細	Goat polyclonal to Ionotropic Glutamate receptor 4
由来種	Goat
特異性	ab77407 is expected to recognise all three reported isoforms (NP_000820.3; NP_001070711.1; NP_001070712.1).
アプリケーション	<b>適用あり:</b> WB, ELISA
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat, Chicken, Cow, Dog, Chimpanzee, Cynomolgus monkey, Rhesus monkey 
免疫原	Synthetic peptide: C-KKLDQREYPGSETP , corresponding to internal sequence amino acids 280-293 of Human Ionotropic Glutamate receptor 4 (NP_000820.3; NP_001070711.1; NP_001070712.1). <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
ポジティブ・コントロール	Human brain (cerebellum) lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 0.5% BSA, Tris saline, pH 7.3
精製度	Immunogen affinity purified
特記事項 (精製)	ab77407 is purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunising peptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab77407** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
----------	-----------	------

WB

ELISA

#### 追加情報

Peptide ELISA: Antibody detection limit dilution 1/128000.

WB: Use at a concentration of 0.3 - 1 µg/ml. Detects a band of approximately 90 kDa (predicted molecular weight: 101 kDa).

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

#### ターゲット情報

##### 機能

Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist.

##### 配列類似性

Belongs to the glutamate-gated ion channel (TC 1.A.10.1) family. GRIA4 subfamily.


##### 翻訳後修飾

Palmitoylated. Depalmitoylated upon glutamate stimulation. Cys-611 palmitoylation leads to Golgi retention and decreased cell surface expression. In contrast, Cys-837 palmitoylation does not affect cell surface expression but regulates stimulation-dependent endocytosis.

##### 細胞内局在

Cell membrane. Cell junction > synapse > postsynaptic cell membrane.

#### 画像



Western blot - Ionotropic Glutamate receptor 4 antibody (ab77407)

Anti-Ionotropic Glutamate receptor 4 antibody (ab77407) at 0.3 µg/ml + human cerebellum lysate in RIPA buffer at 35 µg

**Predicted band size:** 101 kDa

**Observed band size:** 90 kDa

Primary incubation was 1 hour. Detected by chemiluminescence.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors