

Product datasheet

Anti-MERTK antibody ab110108

1 Abreviews 1 References 画像数 1

製品の概要

製品名	Anti-MERTK antibody
製品の詳細	Goat polyclonal to MERTK
アプリケーション	<b>適用あり:</b> ELISA, IHC-P
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide: C-DSSEGSEVLM , corresponding to C terminal amino acids 990-999 of Human MERTK <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
ポジティブ・コントロール	Human Adrenal tissue.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab110108** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		1/2000.
IHC-P		Use a concentration of 3.75 µg/ml.

## ターゲット情報


---

機能	In case of filovirus infection, seems to function as a cell entry factor.
組織特異性	Not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines.
関連疾患	Defects in MERTK are the cause of retinitis pigmentosa type 38 (RP38) [MIM:613862]. RP38 is a retinal dystrophy belonging to the group of pigmentary retinopathies. Retinitis pigmentosa is characterized by retinal pigment deposits visible on fundus examination and primary loss of rod photoreceptor cells followed by secondary loss of cone photoreceptors. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well.
配列類似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. AXL/UFO subfamily. Contains 2 fibronectin type-III domains. Contains 2 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 protein kinase domain.
細胞内局在	Membrane.

---

## 画像

---



ab110108 at 3.75 µg/ml staining MERTK in Formalin-Fixed, Paraffin-Embedded Human Adrenal tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - MERTK antibody (ab110108)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors