


Product datasheet

Anti-MNDA antibody ab118371

画像数 1

製品の概要

製品名	Anti-MNDA antibody
製品の詳細	Mouse monoclonal to MNDA
由来種	Mouse
アプリケーション	適用あり: IHC-P
種交差性	交差種: Human 交差が予測される動物種: Cynomolgus monkey 
免疫原	Synthetic peptide, corresponding to amino acids 311-408 of Human MNDA (NP_002423) with a proprietary tag.
ポジティブ・コントロール	Human kidney tissue
特記事項	Abcam is committed to meeting high standards of ethical manufacturing and as such, we will be discontinuing this product, which has been generated by the ascites method, within the next year. We are sorry for any inconvenience this may cause. If you would like help finding an alternative product, please do not hesitate to contact our scientific support team.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.20 Constituent: 99% PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab118371** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリ ケーショ ン	Abreviews	特記事項
------------------	-----------	------

IHC-P Use a concentration of 3 µg/ml. Antigen retrieval is not essential but may optimise staining. Place sample in 1X citrate buffer (pH 6.0) in pressure cooker under 125°C for 4min and under 90°C for 45min.

ターゲット情報

機能	May act as a transcriptional activator/repressor in the myeloid lineage. Plays a role in the granulocyte/monocyte cell-specific response to interferon. Stimulates the DNA binding of the transcriptional repressor protein YY1.
組織特異性	Expressed constitutively in cells of the myeloid lineage. Found in promyelocyte stage cells as well as in all other stage cells including peripheral blood monocytes and granulocytes. Also appear in myeloblast cells in some cases of acute myeloid Leukemia.
配列類似性	Contains 1 DAPIN domain. Contains 1 HIN-200 domain.
ドメイン	Its N-terminal half (200 amino acids) is sufficient for maximum enhancement of YY1 DNA binding and a portion of this sequence is responsible for binding YY1.
細胞内局在	Nucleus. Cytoplasm. Uniformly distributed throughout the interphase cell nucleus. Associates with chromatin.

画像



Formalin-fixed, paraffin-embedded Human kidney tissue labelled with ab118371 at 3 µg/ml

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MNDA antibody (ab118371)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors