

Product datasheet

Anti-Muscarinic Acetylcholine Receptor M4 antibody - Carboxyterminal end ab41170

1 References

製品の概要

製品名	Anti-Muscarinic Acetylcholine Receptor M4 antibody - Carboxyterminal end
製品の詳細	Rabbit polyclonal to Muscarinic Acetylcholine Receptor M4 - Carboxyterminal end
由来種	Rabbit
特異性	Cross reactivity for ab41170: M4 Muscarinic Receptor (469-478): 100% M4 Muscarinic Receptor: 70% No cross reactivity was observed in: M1 Muscarinic Receptor(451-460) M1 Muscarinic Receptor M2 Muscarinic Receptor(457-466) M2 Muscarinic Receptor M3 Muscarinic Receptor(580-589) M3 Muscarinic Receptor M5 Muscarinic Receptor(519-531) M5 Muscarinic Receptor
アプリケーション	適用あり: ICC, WB, ELISA
種交差性	交差種: Mouse, Rat
免疫原	peptide analogue of the C terminal of rat muscarinic M4 receptor covalently attached onto a carrier protein.
ポジティブ・コントロール	Rat brain tissue, whole brain homogenate and cultured cells.

製品の特性

製品の状態	Lyophilised:Reconstitute the lyophilized pooled antisera with 0.1 ml of 10 mg/ml BSA and 0.01% Sodium Azide in PBS.
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None Constituents: Whole serum
精製度	Whole antiserum
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab41170** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ICC		
WB		
ELISA		

追加情報

ELISA: Use at an assay dependent dilution.
ICC: 1/500.
WB: 1/400. Predicted molecular weight: 53 kDa.
This antibody is very sensitive to blocking agent. We recommend using whole goats serum for optimal results. Avoid any bovine products (BSA and milk).

With this antibody, we have found that blocking with 5% goat or donkey serum significantly reduces background as compared to BSA or milk.

Not yet tested in other applications.
Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性

The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine to these receptors and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The clinical implications of this receptor are unknown; however, mouse studies link its function to adenylate cyclase inhibition. Muscarinic acetylcholine receptor M4 has been reported to be expressed in many regions of the rat brain, particularly the striatum and hippocampus.

細胞内局在

Cell Membrane

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors