

## Product datasheet

# Anti-NGDN antibody ab121124

画像数 3

### 製品の概要

製品名	Anti-NGDN antibody
製品の詳細	Rabbit polyclonal to NGDN
アプリケーション	<b>適用あり:</b> WB, ICC/IF, IHC-P
種交差性	<b>交差種:</b> Human
免疫原	Recombinant fragment, corresponding to amino acids 61-209 of Human NGDN (Q8NEJ9). <a href="#">Run BLAST with ExPASy</a> <a href="#">Run BLAST with NCBI</a>
ポジティブ・コントロール	NGDN transfected HEK293T lysate, Human skin tissue

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab121124** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.

アプリケーション	Abreviews	特記事項
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Use buffer at pH 6 for antigen retrieval.

## ターゲット情報


機能	Involved in the translational repression of cytoplasmic polyadenylation element (CPE)-containing mRNAs.
配列類似性	Belongs to the SAS10 family.
細胞内局在	Nucleus. Cytoplasm. Cell projection > axon. Cell projection > dendrite. Cell projection > filopodium. Detected in axons, dendrites and filopodia. Colocalized with EIF4E in neurites.

## 画像



Immunofluorescent staining of human cell line U-251 MG shows positivity in nucleus, nucleoli & mitochondria.

Immunocytochemistry - Anti-NGDN antibody (ab121124)




Western blot - Anti-NGDN antibody (ab121124)

**All lanes :** Anti-NGDN antibody (ab121124)  
at 1/250 dilution

**Lane 1 :** vector only transfected HEK293T  
lysate

**Lane 2 :** NGDN-transfected HEK293T lysate  
(Co-expressed with a C-terminal myc-DDK  
tag (~3.1 kDa))

Developed using the ECL technique  
10-20%  
SDS PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-  
embedded sections) - Anti-NGDN antibody  
(ab121124)

ab121124 at 1/350 dilution staining NGDN in  
Paraffin-embedded Human skin tissue by  
Immunohistochemistry. Note: strong nuclear  
positivity in epidermal cells.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors