

Product datasheet

Anti-Oncostatin M antibody ab9633

1 References

製品の概要

製品名	Anti-Oncostatin M antibody
製品の詳細	Rabbit polyclonal to Oncostatin M
アプリケーション	<b>適用あり:</b> WB, ELISA, Neutralising
種交差性	<b>交差種:</b> Human
免疫原	Highly pure (>98%) recombinant hOncostatin-M (human Oncostatin-M).

製品の特性

製品の状態	Lyophilised:Reconstitute with 200µl of sterile water.
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	PBS, pH 7.4, no preservative, sterile filtered
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	unknown
軽鎖の種類	unknown

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab9633** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		
ELISA		
Neutralising		

追加情報

Neutralization: To yield one-half maximal inhibition [ND<sub>50</sub>] of the biological activity of hOncostatin-M (10 ng/ml), a concentration of 0.1 - 0.175 µg/ml of this antibody is required

ELISA:To detect hOncostatin-M by direct ELISA (using 100µl/well antibody solution) a

concentration of at least 0.5µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hOncostatin-M.

Western Blot: To detect hOncostatin-M by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hOncostatin-M is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

## ターゲット情報

---

機能	Growth regulator. Inhibits the proliferation of a number of tumor cell lines. Stimulates proliferation of AIDS-KS cells. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. Uses both type I OSM receptor (heterodimers composed of LIPR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST).
配列類似性	Belongs to the LIF/OSM family.
細胞内局在	Secreted.

---

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors