

## Product datasheet

# Anti-PHD1/prolyl hydroxylase antibody ab52726

1 References [画像数 1](#)

### 製品の概要

<b>製品名</b>	Anti-PHD1/prolyl hydroxylase antibody
<b>製品の詳細</b>	Mouse polyclonal to PHD1/prolyl hydroxylase
<b>由来種</b>	Mouse
<b>アプリケーション</b>	<b>適用あり:</b> WB
<b>種交差性</b>	<b>交差種:</b> Caenorhabditis elegans
<b>免疫原</b>	Recombinant fusion protein: RNPHEVMPVF RHRFAITWY MDKSERDKAL AKGKESDAAC, corresponding to amino acids 545-644 of Caenorhabditis elegans PHD1  

### 特記事項

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

### 製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
<b>バッファー</b>	Preservative: None Constituents: 50% Glycerol, Whole serum
<b>精製度</b>	Whole antiserum
<b>一次抗体 備考</b>	This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang <i>et al.</i> <a href="#">PubMed: 1545867</a> ; Chambers and Johnston





Western blot - Anti-PHD1/prolyl hydroxylase antibody (ab52726)

**All lanes :** Anti-PHD1/prolyl hydroxylase antibody (ab52726) at 1/1000 dilution

**Lane 1 :** Total protein extract from E. coli with ~50ng to 100ng of a recombinant fusion protein of an irrelevant antigen

**Lane 2 :** Total protein extract from E. coli with ~50ng to 500ng of the antigen (tagged recombinant fusion protein)

### Secondary

**All lanes :** Rabbit anti-mouse IgG + IgM, horseradish peroxidase conjugated at 1/5000 dilution

**Predicted band size:** 44 kDa

**Observed band size:** 38 kDa

The molecular weight of the band on the western blot does not correspond to the molecular weight of the natural protein because only a fragment of the gene is used.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors