

## Product datasheet

# Anti-Rb antibody [1F8], prediluted ab17119

### 画像数 1

#### 製品の概要

製品名	Anti-Rb antibody [1F8], prediluted
製品の詳細	Mouse monoclonal [1F8] to Rb, prediluted
アプリケーション	適用あり: IHC-P
種交差性	交差種: Human
免疫原	Recombinant full length human Rb protein.
ポジティブ・コントロール	Colon carcinoma

#### 製品の特性

製品の状態	Prediluted
保存方法	Shipped at 4°C. Store at +4°C.
ポリ/モノ	モノクローナル
クローン名	1F8
アイソタイプ	IgG1
軽鎖の種類	kappa

#### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab17119** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
IHC-P		

#### 追加情報

IHC-P: This antibody has been pretitered and ready to use on formalin-fixed paraffin-embedded as well as acetone fixed cryostat tissue sections. No further titration is required.

Formalin-fixed paraffin-embedded tissue sections require high temperature antigen unmasking with 1mM EDTA buffer, pH 8.0 prior to immunostaining.

Not tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.


## ターゲット情報

---

機能	<p>Key regulator of entry into cell division that acts as a tumor suppressor. Promotes G0-G1 transition when phosphorylated by CDK3/cyclin-C. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, KMT5B and KMT5C, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. Mediates transcriptional repression by SMARCA4/BRG1 by recruiting a histone deacetylase (HDAC) complex to the c-FOS promoter. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC1 repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex (By similarity). In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity.</p>
組織特異性	<p>Expressed in the retina.</p>
関連疾患	<p>Childhood cancer retinoblastoma Bladder cancer Osteogenic sarcoma</p>
配列類似性	<p>Belongs to the retinoblastoma protein (RB) family.</p>
ドメイン	<p>The Pocket domain binds to the threonine-phosphorylated domain C, thereby preventing interaction with heterodimeric E2F/DP transcription factor complexes.</p>
翻訳後修飾	<p>Phosphorylated by CDK6 and CDK4, and subsequently by CDK2 at Ser-567 in G1, thereby releasing E2F1 which is then able to activate cell growth. Dephosphorylated at the late M phase. SV40 large T antigen, HPV E7 and adenovirus E1A bind to the underphosphorylated, active form of pRb. Phosphorylation at Thr-821 and Thr-826 promotes interaction between the C-terminal domain C and the Pocket domain, and thereby inhibits interactions with heterodimeric E2F/DP transcription factor complexes. Dephosphorylated at Ser-795 by calcineurin upon calcium stimulation. CDK3/cyclin-C-mediated phosphorylation at Ser-807 and Ser-811 is required for G0-G1 transition. Phosphorylated by CDK1 and CDK2 upon TGFB1-mediated apoptosis.</p> <p>N-terminus is methylated by METTL11A/NTM1 (By similarity). Monomethylation at Lys-810 by SMYD2 enhances phosphorylation at Ser-807 and Ser-811, and promotes cell cycle progression. Monomethylation at Lys-860 by SMYD2 promotes interaction with L3MBTL1. Acetylation at Lys-873 and Lys-874 regulates subcellular localization, at least during keratinocytes differentiation.</p>
細胞内局在	<p>Nucleus.</p>

## 画像

---



Immunohistochemical analysis of formalin-fixed paraffin-embedded human tonsil tissue sections, labelling Retinoblastoma with ab17119.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Rb antibody [1F8], prediluted (ab17119)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors