

## Product datasheet

# Recombinant Human PSIP1 protein ab82129

### 製品の概要

---

製品名	Recombinant Human PSIP1 protein
タンパク質長	Protein fragment

### 製品の詳細

---

由来	Recombinant
由来	Escherichia coli

### アミノ酸配列

生物種	Human
領域	322 to 530

### 特性

---

Our [Abpromise guarantee](#) covers the use of **ab82129** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE
精製度	> 95 % SDS-PAGE.

製品の状態	Liquid
-------	--------

### 前処理および保存

---

保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. Preservative: None Constituents: 20% Glycerol, 20mM Tris Cl, 100mM Potassium chloride, 1mM DTT, 0.2mM EDTA, pH 8.0
------------	--

### 関連情報

---

機能	Transcriptional coactivator involved in neuroepithelial stem cell differentiation and neurogenesis. Involved in particular in lens epithelial cell gene regulation and stress responses. May play an important role in lens epithelial to fiber cell terminal differentiation. May play a protective role
----	---

during stress-induced apoptosis. Isoform 2 is a more general and stronger transcriptional coactivator. Isoform 2 may also act as an adapter to coordinate pre-mRNA splicing. Cellular cofactor for lentiviral integration.

#### 組織特異性

Widely expressed. Expressed at high level in the thymus. Expressed in fetal and adult brain. Expressed in neurons, but not astrocytes. Markedly elevated in fetal as compared to adult brain. In the adult brain, expressed in the subventricular zone (SVZ), in hippocampus, and undetectable elsewhere. In the fetal brain, expressed in the germinal neuroepithelium and cortical plate regions.

#### 関連疾患

Note=A chromosomal aberration involving PSIP1 is associated with pediatric acute myeloid leukemia (AML) with intermediate characteristics between M2-M3 French-American-British (FAB) subtypes. Translocation t(9;11)(p22;p15) with NUP98. The chimeric transcript is an in-frame fusion of NUP98 exon 8 to PSIP1/LEDGF exon 4.

#### 配列類似性

Belongs to the HDGF family.

Contains 1 PWWP domain.

#### ドメイン

Residues 340-417 are necessary and sufficient for the interaction with HIV-1 IN (IBD domain).

#### 翻訳後修飾

Phosphorylated upon DNA damage, probably by ATM or ATR.

#### 細胞内局在

Nucleus. Remains chromatin-associated throughout the cell cycle.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors