

Product datasheet

Recombinant Human SIRT5 protein ab101133

画像数 1

製品の概要

製品名	Recombinant Human SIRT5 protein
タンパク質長	Full length protein

製品の詳細

由来	Recombinant
由来	Escherichia coli
アミノ酸配列	
アクセッション番号	Q9NXA8
生物種	Human
分子量	60 kDa including tags
領域	1 to 310

特性

Our [Abpromise guarantee](#) covers the use of **ab101133** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Western blot SDS-PAGE
精製度	> 95 % SDS-PAGE. The purity was determined to be >95% by densitometry
製品の状態	Liquid


前処理および保存

保存方法および安定性	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles. pH: 7.50 Constituents: 0.307% Glutathione, 0.00174% PMSF, 0.00385% DTT, 0.79% Tris HCl, 0.00292% EDTA, 25% Glycerol, 0.87% Sodium chloride
------------	--

関連情報

機能	NAD-dependent lysine demalonylase, desuccinylase and deglutarylase that specifically removes malonyl, succinyl and glutaryl groups on target proteins (PubMed:21908771, PubMed:22076378, PubMed:24703693). Activates CPS1 and contributes to the regulation of blood ammonia levels during prolonged fasting: acts by mediating desuccinylation and deglutarylation of CPS1, thereby increasing CPS1 activity in response to elevated NAD levels during fasting (PubMed:22076378, PubMed:24703693). Activates SOD1 by mediating its desuccinylation, leading to reduced reactive oxygen species (PubMed:24140062). Modulates ketogenesis through the desuccinylation and activation of HMGCS2 (By similarity). Has weak NAD-dependent protein deacetylase activity; however this activity may not be physiologically relevant in vivo. Can deacetylate cytochrome c (CYCS) and a number of other proteins in vitro such as UOX.
組織特異性	Widely expressed.
配列類似性	Belongs to the sirtuin family. Class III subfamily. Contains 1 deacetylase sirtuin-type domain.
ドメイン	In contrast to class I sirtuins, class III sirtuins have only weak deacetylase activity. Difference in substrate specificity is probably due to a larger hydrophobic pocket with 2 residues (Tyr-102 and Arg-105) that bind to malonylated and succinylated substrates and define the specificity (PubMed:22076378).
細胞内局在	Mitochondrion; Cytoplasm. Mitochondrion and Mitochondrion matrix. Mitochondrion intermembrane space. Cytoplasm, cytosol. Nucleus. Mainly mitochondrial. Also present extramitochondrially: a fraction is present in the cytosol and very small amounts are also detected in the nucleus.

画像



SDS-PAGE showing ab101133 at approximately 60kDa.

SDS-PAGE - Recombinant Human SIRT5 protein
(ab101133)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors