

## Product datasheet

# Recombinant Human UBE2I / UBC9 protein ab3803

画像数 3

### 製品の概要

製品名	Recombinant Human UBE2I / UBC9 protein
タンパク質長	Full length protein

### 製品の詳細

由来	Recombinant
由来	Escherichia coli
アミノ酸配列	
生物種	Human
分子量	18 kDa

### 特性

Our [Abpromise guarantee](#) covers the use of **ab3803** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**生理活性** The final fraction of enzyme contains a single polypeptide band of 18 kDa.

**アプリケーション** Western blot

Functional Studies

**製品の状態** Liquid

### 備考

This product can be used as part of an assay for sumoylation activity. Human Aos 1 + Uba 2 ([ab3804](#)), Ubc 9 ([ab3803](#)) and Sumo 1 ([ab3801](#)) can be used to promote in vitro sumoylation of a sumoylation marker (human Topoisomerase I protein fragment) ([ab3828](#)). The reaction products can be detected using our Sumo 1 ([ab3819](#) and [ab3824](#)) and Topoisomerase I ([ab3825](#)) antibodies. Sumoylation assays are carried out in a final volume of 20µl in reaction conditions (20 mM Hepes pH 7.5, 5mM MgCl<sub>2</sub>, 2mM ATP). Sumoylation Protocol: 1. Prepare a suitable purified substrate protein. (For the control, use 2µl Topoisomerase I marker for each reaction.) 2. In each reaction, add 4µl E2 to substrate first, then 2µl Sumo 1, 2µl 10x reaction buffer, 2µl E1. Finally, add H<sub>2</sub>O to bring up to 20µl. We would recommend adding fresh 2mM ATP to be sure that sufficient energy is supplied. 3. The best reaction concentration of proteins is as following: Aos 1 + Uba 2: 7.5µg/ml. Ubc 9: 50µg/ml. SUMO 1: 50µg/ml. For the control assay we recommend running the assay at 37°C for 30-60 minutes. 4. Detect the reaction

products by Western blot using a suitable antibody. For the control reaction use 1/1000 dilution of the supplied Topoisomerase I antibody. Four sumoylated bands should be seen on the gel for the control reaction. This assay has been shown to work with crude extracts. Be aware that Uba 2 contains his-rich regions which might cross-react with antibodies against the 6x-His epitope tag. During western analysis with anti-6x-His antibodies, Uba 2 at 80 kDa might be shown. The final fraction of enzyme contains a single polypeptide band of 18 kDa.

## 前処理および保存

---

### 保存方法および安定性

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

10 mM Tris-Cl, pH 7.5, 100 mM NaCl, 100 mM imidazole, 0.5 mM PMSF, 1 mM DTT, and 10 % glycerol

## 関連情報

---

### 機能

Accepts the ubiquitin-like proteins SUMO1, SUMO2, SUMO3 and SUMO4 from the UBLE1A-UBLE1B E1 complex and catalyzes their covalent attachment to other proteins with the help of an E3 ligase such as RANBP2 or CBX4. Necessary for sumoylation of FOXL2 and KAT5. Essential for nuclear architecture and chromosome segregation.

### 組織特異性

Expressed in heart, skeletal muscle, pancreas, kidney, liver, lung, placenta and brain. Also expressed in testis and thymus.

### パスウェイ

Protein modification; protein sumoylation.

### 配列類似性


Belongs to the ubiquitin-conjugating enzyme family.

### 細胞内局在

Nucleus. Cytoplasm. Mainly nuclear. In spermatocytes, localizes in synaptonemal complexes. Recruited by BCL11A into the nuclear body.

## 画像

---




Sumoylation assay



Left: Human topo I (S<sup>35</sup>Met-labeled) control.

Right: Sumoylated human topo I (S<sup>35</sup>Met-labeled)

(7.5 µg/ml E1, 50 µg/ml E2, 50 µg/ml sumo1, 37°C 30min).



Extracts prepared from CEF cells not transfected (-) or transfected (+) with wild-type Src were resolved by SDS-PAGE on a 10% Tris-glycine gel and transferred to nitrocellulose. Membranes were blocked with a 5% BSA-TBST buffer overnight at 4°C, then were incubated with 1.0 µg/mL anti-Src pan antibody. After washing, membranes were incubated with goat F(ab')<sub>2</sub> anti-rabbit IgG alkaline phosphatase and the signal was detected using the Tropix WesternStar method.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors