

# Recombinant human IL-16 protein ab9564

### 製品の詳細

製品名	Recombinant human IL-16 protein
エンドトキシン・レベル	< 0.100 Eu/μg
発現系	Escherichia coli
アクセッション番号	<b>Q14005</b>
タンパク質長	Protein fragment
Animal free	No
由来	Recombinant
生物種	Human
配列	PDLNSSTDSAASASAASDVSVESTAEATVCTVTLEKMSAGLG FSLEGGKG SLHGDKPLTINRIFKGAASEQSETVQPGDEILQLGGTAMQGL TRFEAWNI IKALPDGPVTIVIRRKSLQSKETTAAGDS

### 特性

Our **Abpromise guarantee** covers the use of **ab9564** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE Functional Studies
製品の状態	Lyophilized
備考	The biological activity of this product is determined by its ability to chemoattract human CD4 positive T lymphocytes using a concentration range of 1.0-100 ng/ml.

### 前処理および保存

保存方法および安定性	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. pH: 5.50 Constituents: 0.147% Sodium citrate, 0.0077% DTT This product is an active protein and may elicit a biological response in vivo, handle with caution.
再構成	Reconstitute with distilled water to achieve desired concentration.

## 関連情報

---

<b>機能</b>	<p>Interleukin-16 stimulates a migratory response in CD4+ lymphocytes, monocytes, and eosinophils. Primes CD4+ T-cells for IL-2 and IL-15 responsiveness. Also induces T-lymphocyte expression of interleukin 2 receptor. Ligand for CD4.</p> <p>Isoform 1 may act as a scaffolding protein that anchors ion channels in the membrane.</p> <p>Isoform 3 is involved in cell cycle progression in T-cells. Appears to be involved in transcriptional regulation of SKP2 and is probably part of a transcriptional repression complex on the core promoter of the SKP2 gene. May act as a scaffold for GABPB1 (the DNA-binding subunit the GABP transcription factor complex) and HDAC3 thus maintaining transcriptional repression and blocking cell cycle progression in resting T-cells.</p>
<b>組織特異性</b>	<p>Isoform 3 is expressed in hemopoietic tissues, such as resting T-cells, but is undetectable during active T cell proliferation.</p>
<b>配列類似性</b>	<p>Contains 4 PDZ (DHR) domains.</p>
<b>翻訳後修飾</b>	<p>Isoform 3 is synthesized as a chemo-attractant inactive precursor in hemopoietic tissues and is proteolytically cleaved by caspase-3 to yield IL-16.</p>
<b>細胞内局在</b>	<p>Cytoplasm; Cytoplasm. Nucleus and Secreted.</p>

---

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors