

Product datasheet

Recombinant human Serine/threonine-protein kinase 4 ab60337

画像数 2

製品の概要

製品名 Recombinant human Serine/threonine-protein kinase 4
 タンパク質長 Full length protein

法規制情報

カルタヘナ法

製品の詳細

由来 Recombinant
 由来 Baculovirus infected Sf9 cells
 アミノ酸配列
 生物種 Human

特性

Our [Abpromise guarantee](#) covers the use of **ab60337** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション Functional Studies
 SDS-PAGE
 製品の状態 Liquid
 備考 [ab204853](#) (IRS1 peptide) can be utilized as a substrate for assessing kinase activity

前処理および保存

保存方法および安定性 Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.
 Preservative: None
 Constituents: 25% Glycerol, 50mM Tris HCl, 150mM Sodium chloride, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, pH 7.5
 This product is an active protein and may elicit a biological response in vivo, handle with caution.

関連情報

機能

Stress-activated, pro-apoptotic kinase which, following caspase-cleavage, enters the nucleus and induces chromatin condensation followed by internucleosomal DNA fragmentation. Key component of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Phosphorylation of YAP1 by LATS2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. MST1/MST2 are required to repress proliferation of mature hepatocytes, to prevent activation of facultative adult liver stem cells (oval cells), and to inhibit tumor formation (By similarity). Phosphorylates 'Ser-14' of histone H2B (H2BS14ph) during apoptosis. Phosphorylates FOXO3 upon oxidative stress, which results in its nuclear translocation and cell death initiation.

組織特異性

Ubiquitously expressed.

配列類似性

Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily. Contains 1 protein kinase domain. Contains 1 SARAH domain.


翻訳後修飾

Autophosphorylated on serine and threonine residues.

細胞内局在

Cytoplasm. Nucleus. The caspase-cleaved form cycles between the nucleus and cytoplasm.

画像



Sample Kinase Activity Plot.

Functional Studies - Active human Serine/threonine-protein kinase 4 full length protein (ab60337)



ab60337 on SDS-PAGE, MW ~83kDa.

SDS-PAGE - Active human Serine/threonine-protein kinase 4 full length protein (ab60337)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors