

## Product datasheet

# Anti-Rpt5 antibody ab21745

画像数 1

### 製品の概要

<b>製品名</b>	Anti-Rpt5 antibody
<b>製品の詳細</b>	Mouse polyclonal to Rpt5
<b>由来種</b>	Mouse
<b>アプリケーション</b>	<b>適用あり:</b> WB
<b>種交差性</b>	<b>交差種:</b> <i>Saccharomyces cerevisiae</i>
<b>免疫原</b>	Fusion protein: EKIKDNKEKIKNNRQLPYLVANVVEVMDMNEIEDKENSESTTQGGNVNLD NTAVGKAAVVKTSSRQTVFLPMVGLVDPDKLKPNDLVGVNKDSYLILDTL , corresponding to amino acids 60-159 of <i>S. cerevisiae</i> Rpt5. <div style="text-align: right;"> <a href="#">Run BLAST with</a>      <a href="#">Run BLAST with</a> </div>
<b>特記事項</b>	Produced from outbred CD1 mice  This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang <i>et al.</i> <a href="#">PubMed: 1545867</a> ; Chambers and Johnston <a href="#">PubMed: 12910245</a> ; Barry and Johnston <a href="#">PubMed: 9234514</a> ). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an <i>E.coli</i> lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

### 製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>バッファー</b>	Constituents: 50% Glycerol
<b>精製度</b>	Whole antiserum

## 一次抗体 備考

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed: 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

ポリ/モノ

ポリクローナル

アイソタイプ

IgG

## アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab21745** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/1000. Predicted molecular weight: 48 kDa. This antibody has been tested in Western blot against an <i>E.coli</i> lysate containing the partial recombinant fusion protein used as an immunogen. We have no data on detection of endogenous protein.

## ターゲット情報


### 関連性

In the yeast *S. cerevisiae* 12 different members of a novel gene family of putative ATPases have been identified and characterised. Due to their similarities to TBP1, a human immunodeficiency tat binding protein, they have been designated the YTA family or yeast tat binding analogues. All share a region of high similarity of approximately 300 amino acids in length. Rpt5p or YTA1, is one of six ATPases of the 19S regulatory particle of the 26S proteasome involved in the degradation of ubiquitinated substrates.

### 細胞内局在

Nuclear

## 画像



Western blot - Anti-Rpt5 antibody (ab21745)

**All lanes** : Anti-Rpt5 antibody (ab21745) at 1/1000 dilution

**Lane 1** : Total protein extract from E. coli with ~50ng to 100ng of a negative control fusion protein with an irrelevant antigen at 20 ug

**Lane 2** : Total protein extract from E. coli with ~50ng to 500ng of the antigen fusion protein at 20 ug

#### **Secondary**

**All lanes** : Rabbit anti-mouse IgG + IgM, (H+L) horseradish peroxidase conjugated at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size:** 48 kDa

The molecular weight of the band on the western blot does not correspond to the predicted band size above (predicted from the molecular weight of the natural protein) because of the additional mass of the fusion and because the fusion protein only contains a partial fragment of the gene.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### **Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors