

Product datasheet

Anti-SAX antibody ab42105

1 References

製品の概要

製品名	Anti-SAX antibody
製品の詳細	Rabbit polyclonal to SAX
アプリケーション	適用あり: IHC-FoFr, WB, IP, ICC
種交差性	交差種: Drosophila melanogaster
免疫原	Synthetic peptide: GCSTSPDQLPMICSQNSLK conjugated to KLH, corresponding to amino acids 107-125 of Fruit fly (Drosophila melanogaster) SAX

 [Run BLAST with](#)

 [Run BLAST with](#)

法規制情報

医薬用外毒物

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Antibody stabilization buffer, with preservatives, consisting of Hepes, NaCl, KCl, glycerol, BSA and chaotropic agents. Preservative: 0.01% Thimerosal (merthiolate)
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab42105** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
IHC-FoFr		
WB		
IP		
ICC		

追加情報

ICC: 1/200.
 IHC-FoFr: Use at an assay dependent dilution (PMID 19619497).
 IP: 1/200.
 WB: 1/500. Detects a band of approximately 61-65 kDa (predicted molecular weight: 64 kDa).

Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性

SAX (saxophone) protein is a receptor for dpp. In Drosophila, the saxophone gene encodes a type I receptor that mediates signaling by decapentaplegic (dpp), a member of the bone morphogenetic protein (BMP) subgroup of TGF beta-type factors. Over expression or mutation in dpp suppress germline stem cell differentiation. The Saxophone gene is expressed ubiquitously. The Saxophone gene gives two products Brk43E and Berk25 and both gene products interact with TGF super family peptide ligands Dpp. Mutations that completely abolish Saxophone activity cause phenotypes that are similar to partial or complete loss of activity of the dpp ligand. The saxophone products have serine threonine kinase activity responsible for phosphorylation and activation of ligands including pMAD, Screw (Scw), and short gastrulation protein (Sog).

細胞内局在

Membrane; single-pass type I membrane protein

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors