

## Product datasheet

# Anti-STK3 antibody [EP1466Y] ab52641

**KO** 評価済 RabMAb<sup>®</sup>

15 References 画像数 6

### 製品の概要

製品名	Anti-STK3 antibody [EP1466Y]
製品の詳細	Rabbit monoclonal [EP1466Y] to STK3
由来種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-Fr, WB, IP, IHC-P <b>適用なし:</b> Flow Cyt
種交差性	<b>交差種:</b> Mouse, Rat, Human
免疫原	Synthetic peptide within Human STK3 aa 1-100 (N terminal). The exact sequence is proprietary.
ポジティブ・コントロール	ICC/IF: Wildtype HAP1 cells. WB: C6 cell lysate. IHC-P: Human lymphoma tissue.
特記事項	Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>  <b>We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result of this work we are pleased to now offer this antibody in purified format. We are in the process of updating our datasheets. The purified format is designated 'PUR' on our product labels. If you have any questions regarding this update, please contact our Scientific Support team.</b>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP1466Y
アイソタイプ	IgG

## アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab52641** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
IHC-Fr		Use at an assay dependent concentration. PubMed: 24595170
WB		1/10000 - 1/50000. Detects a band of approximately 56 kDa (predicted molecular weight: 56 kDa).
IP		1/50 - 1/100.
IHC-P		1/50 - 1/100. Antigen retrieval step strongly recommended for enhanced signal.

**追加情報** Is unsuitable for Flow Cyt.

## ターゲット情報

<b>機能</b>	Stress-activated, pro-apoptotic kinase which, following caspase-cleavage, enters the nucleus and induces chromatin condensation followed by internucleosomal DNA fragmentation. Key component of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Phosphorylation of YAP1 by LATS2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. MST1/MST2 are required to repress proliferation of mature hepatocytes, to prevent activation of facultative adult liver stem cells (oval cells), and to inhibit tumor formation. Phosphorylates NKX2-1.
<b>組織特異性</b>	Expressed at high levels in adult kidney, skeletal and placenta tissues and at very low levels in adult heart, lung and brain tissues.
<b>配列類似性</b>	Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily. Contains 1 protein kinase domain. Contains 1 SARA domain.
<b>細胞内局在</b>	Cytoplasm. Nucleus. The caspase-cleaved form cycles between nucleus and cytoplasm.


## 画像



Western blot - Anti-STK3 antibody [EP1466Y] (ab52641)


**Lane 1:** Wild-type HAP1 cell lysate (20 µg)  
**Lane 2:** STK3 knockout HAP1 cell lysate (20 µg)  
**Lane 3:** NIH/3T3 cell lysate (20 µg)  
**Lane 4:** 293T cell lysate (20 µg)  
**Lanes 1 - 4:** Merged signal (red and green). Green - ab52641 observed at 56 kDa. Red - loading control, ab8245, observed at 37 kDa.

ab52641 was shown to specifically react with STK3 when STK3 knockout samples were used. Wild-type and STK3 knockout samples were subjected to SDS-PAGE. ab52641 and ab8245 (loading control to GAPDH) were diluted 1/10 000 and 1/2000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-STK3 antibody [EP1466Y] (ab52641)

ab52641 staining STK3 in wild-type HAP1 cells (top panel) and STK3 knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab52641 at 5µg/ml concentration and ab195889 at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.




Immunohistochemical staining of paraffin embedded human breast carcinoma with purified ab52641 at a working dilution of 1/50. The secondary antibody used is [ab97051](#), a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-STK3 antibody [EP1466Y] (ab52641)



ab52641 (purified) at 1/50 immunoprecipitating STK3 in 10  $\mu$ g C6 cell lysate (Lanes 1 and 2, observed at 56 kDa). Lane 3 - Rabbit monoclonal IgG ([ab172730](#)). For western blotting, HRP Veriblot for IP ([ab131366](#)) was used as the secondary antibody (1/10 000). Blocking buffer and concentration: 5% NFDm/TBST Dilution buffer and concentration: 5% NFDm/TBST

Immunoprecipitation - Anti-STK3 antibody [EP1466Y] (ab52641)



Western blot - Anti-STK3 antibody [EP1466Y]  
(ab52641)

**All lanes :** Anti-STK3 antibody [EP1466Y]  
(ab52641) at 1/50000 dilution (purified)

**Lane 1 :** C6 whole cell lysate

**Lane 2 :** NIH/3T3 whole cell lysate

Lysates/proteins at 10 µg per lane.

#### Secondary


**All lanes :** Goat Anti-Rabbit IgG H&L (HRP)  
(ab97051) at 1/50000 dilution

**Predicted band size:** 56 kDa

**Observed band size:** 56 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Western blot - Anti-STK3 antibody [EP1466Y]  
(ab52641)

**All lanes :** Anti-STK3 antibody [EP1466Y]  
(ab52641) at 1/50000 dilution (purified)

**Lane 1 :** HEK293 whole cell lysate

**Lane 2 :** HeLa whole cell lysate

Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP)  
(ab97051) at 1/20000 dilution

**Predicted band size:** 56 kDa

**Observed band size:** 56 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors