abcam

Product datasheet

Anti-Serotonin antibody ab6444

1 Abreviews 1 References

製品の概要

製品名	Anti-Serotonin antibody
製品の詳細	Rabbit polyclonal to Serotonin
由来種	Rabbit
アプリケーション	**適用あり:** ELISA, ICC
種交差性	**交差種:** Species independent
免疫原	Chemical/ Small Molecule corresponding to Serotonin. Synthetic 5-Hydroxytryptamine conjugated to protein carrier (PC)
特記事項	ab6444 specific to conjugated 5-Hydroxytryptamine (Serotonin) The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
精製度	Ammonium Sulphate Precipitation
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee	**Abpromise保証は、次のテスト済みアプリケーションにおけるab6444の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent concentration.
ICC		Use at an assay dependent concentration.

ターゲット情報

関連性

Serotonin (5-hydroxytryptamine, or 5-HT) is a monoamine neurotransmitter synthesised in serotonergic neurons in the central nervous system and enterochromaffin cells in the gastrointestinal tract. Serotonin plays an important part in the biochemistry of depression, migraine, bipolar disorder and anxiety. It is also believed to be influential on sexuality and appetite. 5-HT is generally thought not to be released from synaptic terminal buttons in the manner of classical neurotransmission but from serotonergic varacosities into the extra neuronal space. From here it is free to diffuse over a relatively large region of space (>20µm) and activate 5-HT receptors located on the dendrites, cell bodies and presynaptic terminals of adjacent neurons. Serotonergic action is terminated primarily via uptake of 5-HT from the synapse. This is through the specific monoamine transporter for 5-HT, 5-HT reuptake transporter, on the presynaptic neuron. The pharmacology of 5-HT is extremely complex, with its actions being mediated by a large and diverse range of 5-HT receptors.

細胞内局在

Cytoplasmic and Secreted

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit **https://www.abcam.co.jp/abpromise** or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors