

Product datasheet

Anti-TNF Receptor II antibody ab74315

★★★★☆ 1 Abreviews [画像数 2](#)

製品の概要

製品名	Anti-TNF Receptor II antibody
製品の詳細	Rabbit polyclonal to TNF Receptor II
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide conjugated to KLH derived from within residues 400 to the C-terminus of Human TNF Receptor II. Immunogenの所有権に関して (Peptide available as ab74611 .)
ポジティブ・コントロール	This antibody gave a positive signal in the following whole cell lysates: Caco 2, HEK293

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab74315** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 48 kDa (predicted molecular weight: 48 kDa).
IHC-P	★★★★☆	Use at an assay dependent concentration.

ターゲット情報

機能	Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.
配列類似性	Contains 4 TNFR-Cys repeats.
翻訳後修飾	Phosphorylated; mainly on serine residues and with a very low level on threonine residues. A soluble form (tumor necrosis factor binding protein 2) is produced from the membrane form by proteolytic processing.
細胞内局在	Secreted and Cell membrane.

画像



All lanes : Anti-TNF Receptor II antibody (ab74315) at 1 µg/ml

Lane 1 : Caco 2 (Human colonic carcinoma cell line) Whole Cell Lysate (ab76828)

Lane 2 : HEK293 (Human) Whole Cell Lysate (ab52256)

Lysates/proteins at 10 µg per lane.

Secondary

Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) (ab65484) at 1/3000 dilution


Performed under reducing conditions.

Predicted band size : 48 kDa

Observed band size : 52 kDa

Additional bands at : 70 kDa. We are unsure as to the identity of these extra bands.

Exposure time : 20 minutes



Western blot - TNF Receptor II antibody (ab74315)

All lanes : Anti-TNF Receptor II antibody (ab74315) at 1 µg/ml

Lane 1 : Heart (Rat) Tissue Lysate

Lane 2 : Thymus (Rat) Tissue Lysate

Lysates/proteins at 10 µg per lane.

Secondary

Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (ab97080) at 1/5000 dilution

Developed using the ECL technique

Performed under reducing conditions.

Predicted band size : 48 kDa

Observed band size : 75 kDa

Additional bands at : 50 kDa (possible cleavage fragment).

Exposure time : 8 minutes

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors