

Product datasheet

Anti-Tissue Factor (phospho S290) antibody ab62251

★★★★☆ 1 Abreviews 1 References 画像数 2

製品の概要

製品名	Anti-Tissue Factor (phospho S290) antibody
製品の詳細	Rabbit polyclonal to Tissue Factor (phospho S290)
特異性	ab62251 detects endogenous levels of Tissue Factor only when phosphorylated at serine 290.
アプリケーション	適用あり: ELISA, IHC-P, WB
種交差性	交差種: Human
免疫原	Synthetic phosphopeptide derived from human Tissue Factor around the phosphorylation site of serine 290 (E-N-S ^P -P-L).
ポジティブ・コントロール	Human brain tissue.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 50% Glycerol, PBS, 150mM Sodium chloride, pH 7.4
精製度	Immunogen affinity purified
特記事項(精製)	ab62251 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab62251** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
ELISA		1/10000.

アプリケーション	Abreviews	特記事項
IHC-P		1/50 - 1/100.
WB	★★★★☆	1/2000. PubMed: 21310953


ターゲット情報

機能	Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited proteolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.
配列類似性	Belongs to the tissue factor family.
細胞内局在	Membrane.

画像



ab62251 at 1/50 - 1/100 dilution staining Tissue Factor in Human brain by Immunohistochemistry, Paraffin-embedded tissue, in the absence or presence of the immunising peptide.



Anti-Tissue Factor (phospho S290) antibody (ab62251) at 1/2000 dilution + whole cell lysate prepared from transfected human endothelial cells expressing the human full length Tissue Factor (phospho S290) (treated with PAR2-agonist peptide for 90 minutes) at 20 µg/ml

Secondary

Goat anti-rabbit IgG-HRP at 1/1000 dilution

Observed band size : 46 kDa

Image courtesy of Dr C Ettelaie by Abreview.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors