

Product datasheet

Anti-Transferrin antibody [11D3] (HRP) ab10208

★★★★☆ 1 Abreviews 3 References 画像数 1

製品の概要

製品名	Anti-Transferrin antibody [11D3] (HRP)
製品の詳細	Mouse monoclonal [11D3] to Transferrin (HRP)
由来種	Mouse
標識	HRP
アプリケーション	適用あり: ELISA, Sandwich ELISA, WB
種交差性	交差種: Human
免疫原	Placental transferrin receptor or transferrin.
ポジティブ・コントロール	human brain
特記事項	Concentration varies from lot to lot and can be provided on request. Abcam is committed to meeting high standards of ethical manufacturing and has decided to discontinue this product by June 2019 as it has been generated by the ascites method. We are sorry for any inconvenience this may cause.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	Preservative: 0.05% Proclin Constituents: PBS, pH 7.4
精製度	Protein A purified
特記事項(精製)	Purified by chromatography on protein A sepharose. Purity is tested by electrophoresis.
ポリ/モノ	モノクローナル
クローン名	11D3
ミエローマ	Sp2/0
アイソタイプ	IgG1

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab10208** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent dilution.
Sandwich ELISA		Use at an assay dependent dilution. Can be used as both coating and conjugate antibody when paired with recommended pair for detection of transferrin.
WB	★★★★☆	1/500. Detects a band of approximately 77 kDa (predicted molecular weight: 77 kDa).

ターゲット情報

機能	Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation.
組織特異性	Expressed by the liver and secreted in plasma.
関連疾患	Defects in TF are the cause of atransferrinemia (ATRAF) [MIM:209300]. Atransferrinemia is rare autosomal recessive disorder characterized by iron overload and hypochromic anemia.
配列類似性	Belongs to the transferrin family. Contains 2 transferrin-like domains.
細胞内局在	Secreted.

画像



Western blot - Anti-Transferrin antibody [11D3]
(HRP) (ab10208)

This image is courtesy of Ruma Raha-Chowdhury,
University Of Cambridge, United Kingdom

All lanes : Anti-Transferrin antibody [11D3]
(HRP) (ab10208) at 1/500 dilution

All lanes : human brain

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : DAKO Cytomation

Conjugation: HRP at 1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 77 kDa

Observed band size: 77 kDa

Additional bands at: many <77 kDa

(possible non-specific binding)

Exposure time: 1 minute

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors