

## Product datasheet

# Anti-V5 tag antibody ab9113

★★★★★ 1 Abreviews 3 References 画像数 1

### 製品の概要

製品名	Anti-V5 tag antibody
製品の詳細	Chicken polyclonal to V5 tag
由来種	Chicken
特異性	Antibody specificity was verified by ELISA against the peptide conjugated to BSA (GKPIPPLLGLDST-BSA). A 1:100,000 dilution of the antibody gave an O.D.=1.0 in a 15 minute reaction using HRP conjugated goat anti chicken IgY (IgG) and TMB as the substrate. Appropriate specificity controls were run.
アプリケーション	<b>適用あり:</b> ELISA, WB, ICC/IF, ICC
免疫原	GKPIPPLLGLDST (V5 epitope) conjugated with KLH.

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	PBS, pH 7.2, sodium azide, 0.1%
精製度	IgY fraction
特記事項(精製)	Antibodies were immunoaffinity purified using the peptide immobilized on a solid support. Antibody concentration was determined by extinction coefficient : O.D. 1.32 at 280nm equals 1.0 mg of IgY (IgG).
ポリ/モノ	ポリクローナル
アイソタイプ	unknown
軽鎖の種類	unknown

### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab9113** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
ELISA		Use at an assay dependent dilution.

アプリケーション	Abreviews	特記事項
WB		1/5000. Can be blocked with <a href="#">V5 tag peptide (ab15829)</a> .
ICC/IF	★★★★★	Use at an assay dependent dilution.
ICC		Use at an assay dependent dilution.

## ターゲット情報

**関連性** The V5 epitope tag is derived from a small epitope (Pk) present on the P and V proteins of the paramyxovirus of simian virus 5 (SV5). The V5 tag is usually used with all 14 amino acids (GKPIPPELLGLDST), although it has also been used with a shorter 9 amino acid sequence (IPNPLLGLD).

## 画像



Immunocytochemistry/ Immunofluorescence - Anti-V5 tag antibody (ab9113)

This image is courtesy of an anonymous Abreview

ab9113 staining human fibrosarcoma cells (HT1080) by ICC/IF. Cells were transfected with V5- tagged transmembrane protein and PFA fixed without undergoing permeabilization and blocking steps. The primary antibody was used at 10µg/ml and incubated with the sample for 1 hour at 20°C. An Alexa Fluor® 546 conjugated donkey polyclonal anti-chicken (H+L) antibody, diluted 1/500, was used as the secondary.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors