

Product datasheet

Anti-ZIC5 antibody ab115566

画像数 3

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-ZIC5 antibody |
| 製品の詳細 | Goat polyclonal to ZIC5 |
| アプリケーション | 適用あり: ELISA, IHC-P |
| 種交差性 | 交差種: Human |
| 免疫原 | Synthetic peptide: EIYGNPEVVRTIH with a Cysteine residue linker, corresponding to C terminal amino acids 651-663 of Human ZIC5 (NP_149123.2). Run BLAST with Run BLAST with |
| ポジティブ・コントロール | Human Prostate, Human Skin and Human Small Intestine tissues. |

製品の特性

| | |
|--------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles. |
| バッファー | pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA |
| 精製度 | Protein G purified |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab115566** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|-----------|
| ELISA | | 1/128000. |

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| IHC-P | | Use a concentration of 3.75 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |


ターゲット情報

| | |
|--------------|--|
| 機能 | Essential for neural crest development, converting cells from an epidermal fate to a neural crest cell fate. Binds to DNA. |
| 配列類似性 | Belongs to the GLI C2H2-type zinc-finger protein family. Contains 4 C2H2-type zinc fingers. |
| 細胞内局在 | Nucleus. |


画像



ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Prostate tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.



ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Skin tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.



ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Small intestine tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen. .

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ZIC5 antibody (ab115566)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors