

Product datasheet

Anti-Transferrin Receptor antibody [2B6] (HRP) ab10252

製品の概要

製品名	Anti-Transferrin Receptor antibody [2B6] (HRP)
製品の詳細	Mouse monoclonal [2B6] to Transferrin Receptor (HRP)
由来種	Mouse
標識	HRP
特異性	This antibody reacts with human soluble transferrin receptor.
アプリケーション	適用あり: Sandwich ELISA, ELISA
種交差性	交差種: Human
免疫原	Recombinant full length protein (Human).
特記事項	Concentration varies from lot to lot and can be provided on request.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	Preservative: 0.05% Proclin Constituents: PBS, pH 7.4
精製度	Protein A purified
特記事項 (精製)	Purity tested by electrophoresis.
ポリ/モノ	モノクローナル
クローン名	2B6
ミエローマ	Sp2/0
アイソタイプ	IgG2a

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab10252** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
----------	-----------	------

Sandwich ELISA

ELISA

追加情報

ELISA: Use at an assay dependent dilution.
 sELISA: Use at an assay dependent dilution.
 Can be paired for ELISA with [Mouse monoclonal \[11F5\] to Transferrin Receptor \(ab10249\)](#) and [Mouse monoclonal \[5B11\] to Transferrin Receptor \(ab10253\)](#).
 Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能 Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake (PubMed:26642240).
 (Microbial infection) Acts as a receptor for new-world arenaviruses: Guanarito, Junin and Machupo virus.

関連疾患 Immunodeficiency 46

配列類似性 Belongs to the peptidase M28 family. M28B subfamily.
 Contains 1 PA (protease associated) domain.

翻訳後修飾 N- and O-glycosylated, phosphorylated and palmitoylated. The serum form is only glycosylated. Proteolytically cleaved on Arg-100 to produce the soluble serum form (sTfR).
 Palmitoylated on both Cys-62 and Cys-67. Cys-62 seems to be the major site of palmitoylation.

細胞内局在 Secreted and Cell membrane. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors