

Product datasheet

Anti-BAIAP2 antibody ab110693

画像数 1

製品の概要

製品名	Anti-BAIAP2 antibody
製品の詳細	Rabbit polyclonal to BAIAP2
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Mouse 交差が予測される動物種: Rat, Horse, Chicken, Guinea pig, Cow, Cat, Dog
免疫原	Synthetic peptide corresponding to a region within N terminal amino acids 101-150 (LLTQLEQKVE LDSRYLSAAL KKYQTEQRSK GDALDKCQAE LKKLRKKSQG) of Mouse BAIAP2 (NP_001032844). Run BLAST with ExPASy Run BLAST with NCBI
ポジティブ・コントロール	Mouse Heart lysate

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	Constituents: PBS, 2% Sucrose
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab110693** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 59 kDa. Good results were obtained when blocked with 5% non-fat dry milk in 0.05% PBS-T.

ターゲット情報

機能	Adapter protein that links membrane-bound small G-proteins to cytoplasmic effector proteins. Necessary for CDC42-mediated reorganization of the actin cytoskeleton and for RAC1-mediated membrane ruffling. Involved in the regulation of the actin cytoskeleton by WASF family members and the Arp2/3 complex. Plays a role in neurite growth. Acts synergetically with ENAH to promote filipodia formation.
組織特異性	Isoform 1 and isoform 4 are expressed almost exclusively in brain. Isoform 4 is barely detectable in placenta, prostate and testis. A short isoform is ubiquitous, with the highest expression in liver, prostate, testis and placenta.
配列類似性	Contains 1 IMD (IRSp53/MIM homology) domain. Contains 1 SH3 domain.
ドメイン	The IMD domain forms a coiled coil. The isolated domain can induce actin bundling and filopodia formation. In the absence of G-proteins intramolecular interaction between the IMD and the SH3 domain gives rise to an auto-inhibited state of the protein. Interaction of the IMD with RAC1 or CDC42 leads to activation. The SH3 domain interacts with ATN1, BAI1, WASF1, WASF2, SHANK1, DIAPH1 and ENAH.
翻訳後修飾	Phosphorylated on tyrosine residues by INSR in response to insulin treatment.
細胞内局在	Cytoplasm. Membrane. Cell projection > filopodium. Cell projection > ruffle. Detected throughout the cytoplasm in the absence of specific binding partners. Detected in filopodia and close to membrane ruffles.

画像



Anti-BAIAP2 antibody (ab110693) at 1 µg/ml
+ Mouse Heart lysate at 10 µg

Predicted band size: 59 kDa

Gel concentration: 12%

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors