


Product datasheet

Anti-SIRT6 antibody ab118487

画像数 3

製品の概要

製品名	Anti-SIRT6 antibody
製品の詳細	Rabbit polyclonal to SIRT6
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Dog 
免疫原	Synthetic peptide conjugated to KLH corresponding to residues in Human SIRT6 (NP_057623.1).
ポジティブ・コントロール	Human testis tissue; HepG2 cell lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C.
バッファー	Constituents: 2% Sucrose, 97% PBS
精製度	Protein A purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab118487** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1.25 µg/ml. Predicted molecular weight: 39 kDa.
IHC-P		Use a concentration of 10 µg/ml.

ターゲット情報

機能

NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Acts as a corepressor of the transcription factor HIF1A to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Regulates the production of TNF protein. Has a role in the regulation of life span (By similarity). Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD(+) in the absence of acetylated substrate.


配列類似性

Belongs to the sirtuin family. Class IV subfamily.
Contains 1 deacetylase sirtuin-type domain.

細胞内局在


Nucleus, nucleoplasm. Predominantly nuclear. Associated with telomeric heterochromatin regions.

Anti-SIRT6 antibody 画像




Anti-SIRT6 antibody (ab118487) + U2OS cell lysate

Predicted band size : 39 kDa



ab118487 at 10µg/ml staining SIRT6 in Formalin-fixed, Paraffin-embedded Human testis tissue by immunohistochemistry.



Anti-SIRT6 antibody (ab118487) at 1.25 μ g/ml
+ HepG2 cell lysate

Predicted band size : 39 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors