

Product datasheet

Anti-HAUSP / USP7 antibody [3D3] ab119364

KO 評価済

画像数 7

製品の概要

製品名	Anti-HAUSP / USP7 antibody [3D3]
製品の詳細	Mouse monoclonal [3D3] to HAUSP / USP7
由来種	Mouse
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	Recombinant full length Human HAUSP/USP7 produced in HEK293T cells (NP_003461).
ポジティブ・コントロール	HEK293T cell lysate transfected with pCMV6-ENTRY HAUSP/USP7 cDNA; Human tonsil, thyroid carcinoma, endometrium, endometrium adenocarcinoma and bladder carcinoma tissue.
特記事項	Dilute in PBS (pH7.3) before use. Stable for 12 months from date of receipt.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 48% PBS, 50% Glycerol, 1% BSA
精製度	Protein G purified
特記事項 (精製)	ab119364 is purified from Mouse ascites fluid by affinity chromatography.
ポリモノ	モノクローナル
クローン名	3D3
アイソタイプ	IgG1

アプリケーション


Our [Abpromise guarantee](#) covers the use of **ab119364** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/2000. Predicted molecular weight: 128 kDa.
IHC-P		1/150.

ターゲット情報

機能	<p>Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation. Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:25283148). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis. Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity. In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML. Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6. Contributes to the overall stabilization and trans-activation capability of the herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110 during HSV-1 infection. Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:21745816). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains. Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:23973222).</p>
組織特異性	Widely expressed. Overexpressed in prostate cancer.
配列類似性	<p>Belongs to the peptidase C19 family.</p> <p>Contains 1 MATH domain.</p> <p>Contains 1 USP domain.</p>
ドメイン	The C-terminus plays a role in its oligomerization.
翻訳後修飾	<p>Isoform 1: Phosphorylated. Isoform 1 is phosphorylated at positions Ser-18 and Ser-963.</p> <p>Isoform 2: Not phosphorylated.</p> <p>Isoform 1: Polyneddylated. Isoform 2: Not Polyneddylated.</p> <p>Isoform 1 and isoform 2: Not sumoylated.</p> <p>Isoform 1 and isoform 2: Polyubiquitinated by herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110; leading to its subsequent proteasomal degradation. Isoform 1: Ubiquitinated at Lys-869.</p>
細胞内局在	<p>Nucleus. Cytoplasm. Nucleus, PML body. Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies.</p>



Western blot - Anti-HAUSP / USP7 antibody [3D3] (ab119364)

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: HAUSP/USP7 knockout HAP1 cell lysate (20 µg)


Lane 3: HeLa cell lysate (20 µg)

Lane 4: MCF7 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green).

Green - ab119364 observed at 115 kDa. Red - loading control, [ab181602](#), observed at 37 kDa.

ab119364 was shown to recognize HAUSP/USP7 when HAUSP/USP7 knockout samples were used, along with additional cross-reactive bands. Wild-type and HAUSP/USP7 knockout samples were subjected to SDS-PAGE. ab119364 and [ab181602](#) (loading control to GAPDH) were both diluted 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed [ab216772](#) and Goat Anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed [ab216777](#) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-HAUSP / USP7 antibody [3D3]
(ab119364)

All lanes : Anti-HAUSP / USP7 antibody
[3D3] (ab119364) at 1/500 dilution


Lane 1 : HEK293T cell lysate transfected with
pCMV6-ENTRY control cDNA

Lane 2 : HEK293T cell lysate transfected with
pCMV6-ENTRY HAUSP/USP7 cDNA

Lysates/proteins at 5 µg per lane.


Predicted band size: 128 kDa

HEK293T cell lysates were generated from
transient transfection of the cDNA clone
(RC213986)




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-HAUSP / USP7 antibody
[3D3] (ab119364)

ab119364 at 1/150 dilution staining
HAUSP/USP7 in Paraffin-embedded Human
thyroid carcinoma tissue by
Immunohistochemistry.




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-HAUSP / USP7 antibody
[3D3] (ab119364)

ab119364 at 1/150 dilution staining
HAUSP/USP7 in Paraffin-embedded Human
tonsil tissue by Immunohistochemistry.




ab119364 at 1/150 dilution staining
HAUSP/USP7 in Paraffin-embedded Human
bladder carcinoma tissue by
Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-HAUSP / USP7 antibody
[3D3] (ab119364)



ab119364 at 1/150 dilution staining
HAUSP/USP7 in Paraffin-embedded Human
endometrium tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-HAUSP / USP7 antibody
[3D3] (ab119364)



ab119364 at 1/150 dilution staining
HAUSP/USP7 in Paraffin-embedded Human
endometrium adenocarcinoma tissue by
Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-HAUSP / USP7 antibody
[3D3] (ab119364)

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors