


Product datasheet

Anti-iNOS antibody ab21775

1 Abreviews 4 References 画像数 1

製品の概要

<b>製品名</b>	Anti-iNOS antibody
<b>製品の詳細</b>	Mouse polyclonal to iNOS
<b>アプリケーション</b>	<b>適用あり:</b> WB
<b>種交差性</b>	<b>交差種:</b> Rat <b>交差が予測される動物種:</b> Mouse, Rabbit, Human 
<b>免疫原</b>	Fusion protein: KPRYYSISSSQDHTPSEVHLTVAVVTYRTRDGGPLHHGVCSTWINNLKP EDPVPCFVRSVSGFQLPEDPSQPCILIGPGTGIAPFRSFWQQRLHDSQHR , corresponding to amino acids 901-1000 of Rat iNOS.  <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
<b>特記事項</b>	Produced from outbred CD1 mice

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed: 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>バッファー</b>	Constituents: 50% Glycerol
<b>精製度</b>	Whole antiserum

## 一次抗体 備考

This antibody was raised by a genetic immunization technique. Genetic immunization can be used to generate antibodies by directly delivering antigen-coding DNA into the animal, rather than injecting a protein or peptide (Tang *et al.* [PubMed: 1545867](#); Chambers and Johnston [PubMed: 12910245](#); Barry and Johnston [PubMed: 9234514](#)). The animal's cells produce the protein, which stimulates the animal's immune system to produce antibodies against that particular protein. A vector coding for a partial fusion protein was used for genetic immunisation of a mouse and the resulting serum was tested in Western blot against an *E.coli* lysate containing that partial fusion protein. Genetic immunization offers enormous advantages over the traditional protein-based immunization method. DNA is faster, cheaper and easier to produce and can be produced by standard techniques readily amenable to automation. Furthermore, the antibodies generated by genetic immunization are usually of superior quality with regard to specificity, affinity and recognizing the native protein.

ポリ/モノ

ポリクローナル

アイソタイプ

IgG

## アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab21775** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/1000. Predicted molecular weight: 131 kDa. This antibody has been tested in Western blot against an <i>E.coli</i> lysate containing the partial recombinant fusion protein used as an immunogen. We have no data on detection of endogenous protein.

## ターゲット情報

機能	Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such COX2.
組織特異性	Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets.
配列類似性	Belongs to the NOS family. Contains 1 FAD-binding FR-type domain. Contains 1 flavodoxin-like domain.

## 画像



**All lanes** : Anti-iNOS antibody (ab21775) at 1/1000 dilution

**Lane 1** : Total protein extract from E. coli with ~50ng to 100ng of a negative control fusion protein with an irrelevant antigen at 20 ug

**Lane 2** : Total protein extract from E. coli with ~50ng to 500ng of the antigen fusion protein at 20 ug

### Secondary

Rabbit anti-mouse IgG + IgM, (H+L)  
horseradish peroxidase conjugated at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size** : 131 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors