

Product datasheet

Anti-BDNF antibody ab24491

1 References

製品の概要

製品名	Anti-BDNF antibody
製品の詳細	Sheep polyclonal to BDNF
由来種	Sheep
特異性	Less than 1% cross reactivity against mouse NGF, recombinant human NT3 or Neurotrophin 4.
アプリケーション	適用あり: IHC-FoFr, WB, ICC/IF, Inhibition Assay
種交差性	交差種: Rat, Human
免疫原	Recombinant full length protein (Human)
ポジティブ・コントロール	Brain tissue lysate
特記事項	Antibacterial agents may be required for short term storage. For BDNF, multiple WB bands are possible and expected. The human protein has 5 isoforms (precursors: 28 – 37 kDa) and can be glycosylated (Uniprot: http://www.uniprot.org/uniprot/P23560). The mature form is expected at ~14 kDa (monomer) and the dimer at ~28 kDa.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: None Constituents: Whole serum
精製度	Whole antiserum
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab24491** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
IHC-FoFr		Use at an assay dependent concentration. PubMed: 21667226
WB		1/200 - 1/2000. Predicted molecular weight: 28 kDa.
ICC/IF		Use at an assay dependent dilution.
Inhibition Assay		1/10 - 1/50.

ターゲット情報

機能	During development, promotes the survival and differentiation of selected neuronal populations of the peripheral and central nervous systems. Participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. Major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability.
組織特異性	Brain. Highly expressed in hippocampus, amygdala, cerebral cortex and cerebellum. Also expressed in heart, lung, skeletal muscle, testis, prostate and placenta.
関連疾患	Bulimia nervosa 2 Congenital central hypoventilation syndrome
配列類似性	Belongs to the NGF-beta family.
翻訳後修飾	The propeptide is N-glycosylated and glycosulfated. Converted into mature BDNF by plasmin (PLG).
細胞内局在	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors