

Product datasheet

Anti-beta 2 Microglobulin antibody [F21-21] (Biotin) ab25160

製品の概要

製品名	Anti-beta 2 Microglobulin antibody [F21-21] (Biotin)
製品の詳細	Mouse monoclonal [F21-21] to beta 2 Microglobulin (Biotin)
由来種	Mouse
標識	Biotin
特異性	ab25160 recognises the β 2-microglobulin subunit of MHC Class I molecules
アプリケーション	適用あり: Other, Flow Cyt, IP, WB
種交差性	交差種: Chicken, Turkey
免疫原	The details of the immunogen for this antibody are not available.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.1% Sodium Azide Constituents: PBS
精製度	IgG fraction
ポリ/モノ	モノクローナル
クローン名	F21-21
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab25160** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
Other		

アプリケーション	Abreviews	特記事項
Flow Cyt		
IP		
WB		
AP		
追加情報	<p>AP: Use at an assay dependent dilution. Flow Cyt: Use 1µg for 10⁶ cells. IP: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Predicted molecular weight: 13 kDa.</p> <p>Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user.</p>	
ターゲット情報		
機能	<p>Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system.</p>	
関連疾患	<p>Defects in B2M are the cause of hypercatabolic hypoproteinemia (HYCATHYP) [MIM:241600]. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation.</p> <p>Note=Beta-2-microglobulin may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. Persistently high beta(2)-microglobulin serum levels lead to amyloidosis in patients on long-term hemodialysis.</p>	
配列類似性	<p>Belongs to the beta-2-microglobulin family. Contains 1 Ig-like C1-type (immunoglobulin-like) domain.</p>	
翻訳後修飾	<p>Glycation of Ile-21 is observed in long-term hemodialysis patients.</p>	
細胞内局在	<p>Secreted. Detected in serum and urine.</p>	

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors