


## Product datasheet

# Anti-ErbB 2 antibody - BSA and Azide free ab31836

### 1 References

#### 製品の概要

<b>製品名</b>	Anti-ErbB 2 antibody - BSA and Azide free
<b>製品の詳細</b>	Rabbit polyclonal to ErbB 2 - BSA and Azide free
<b>特異性</b>	ab31836 recognize wild and mutant (oncogenic) forms of c-erbB-2 protein (185kD). Does not crossreact with c-erbB-11 (EGF-R), c-erbB-3,2 or c-erbB-4
<b>アプリケーション</b>	<b>適用あり:</b> WB, IP, IHC-P
<b>種交差性</b>	<b>交差種:</b> Mouse, Rat, Human, Monkey <b>交差が予測される動物種:</b> Cat, Dog 
<b>免疫原</b>	Synthetic peptide: AENPEYLGLDVPV conjugated to KLH, corresponding to C terminal amino acids 1243/1255 of Human c-ErbB2 <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
<b>ポジティブ・コントロール</b>	T47D cells, SKBR3 cells.

#### 製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
<b>バッファー</b>	Preservative: None Constituents: 40% Glycerol, PBS, pH 7.4
<b>精製度</b>	Protein A purified
<b>ポリモノ</b>	ポリクローナル
<b>アイソタイプ</b>	IgG

#### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab31836** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		
IP		
IHC-P		
<b>追加情報</b>	<p>IHC-P: Use at a concentration of 2 - 5 µg/ml for 30 minutes at RT. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. In particular, boil tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes.</p> <p>IP: Use at a concentration of 10 µg/mg protein lysate (Use protein A).</p> <p>WB: Use at a concentration of 5 - 10 µg/ml for 2 hours at RT. Predicted molecular weight: 138 kDa.</p> <p>Not yet tested in other applications. Optimal dilutions/concentrations should be determined by the end user.</p>	
<b>ターゲット情報</b>		
<b>機能</b>	<p>Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.</p>	
<b>組織特異性</b>	<p>Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.</p>	
<b>関連疾患</b>	<p>Hereditary diffuse gastric cancer Glioma Ovarian cancer Lung cancer Gastric cancer</p> <p>Chromosomal aberrations involving ERBB2 may be a cause gastric cancer. Deletions within 17q12 region producing fusion transcripts with CDK12, leading to CDK12-ERBB2 fusion leading to truncated CDK12 protein not in-frame with ERBB2.</p>	
<b>配列類似性</b>	<p>Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily. Contains 1 protein kinase domain.</p>	
<b>翻訳後修飾</b>	<p>Autophosphorylated. Autophosphorylation occurs in trans, i.e. one subunit of the dimeric receptor phosphorylates tyrosine residues on the other subunit (Probable). Ligand-binding</p>	

increases phosphorylation on tyrosine residues (PubMed:27134172). Signaling via SEMA4C promotes phosphorylation at Tyr-1248 (PubMed:17554007). Dephosphorylated by PTPN12 (PubMed:27134172).

#### 細胞内局在

Cytoplasm. Nucleus and Cell membrane. Cytoplasm, perinuclear region. Nucleus. Translocation to the nucleus requires endocytosis, probably endosomal sorting and is mediated by importin beta-1/KPNB1.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors