

Product datasheet

Human breast tissue lysate (ductal carcinoma) - insoluble fraction (female, 45 years) ab43354

製品の概要

<b>製品名</b>	Human breast tissue lysate (ductal carcinoma) - insoluble fraction (female, 45 years)
<b>特記事項</b>	<p>Tissue: Breast Location: Left breast. Diagnosis: Infiltrating ductal carcinoma. Stage: IV T<sub>4</sub>N<sub>2</sub>M<sub>x</sub> Grade: II Sex: Female Age: 45 years Gross findings: Tumor size 8 cm. Cut section hard and yellow. Ipsilateral axillary lymph nodes examined: 8 Microscopic findings: Metastasis to lymph nodes : Positive Vascular infiltration: Positive Tumor necrosis: Positive Cellular differentiation: 2 Nuclear pleomorphism: 2 Mitotic activity: 1 Lymphocytic response: Positive Insoluble protein fraction</p> <p>This product belongs to a range of individual sets of lysates that were extracted from very specific tissues and are gender-, cancer subtype-, tumor grade- and cellular morphology-specific. They allow for consistency, reproducibility, and comparative studies from experiment to experiment, due to the existence of specific (matched) lysates that can be used as controls. Two extractions from each tissue were performed, extraction 1 yielding the soluble protein fraction and extraction 2 the insoluble protein fraction. Please find below a range of products that relate to ab43354: Extraction 1: soluble protein fraction <a href="#">ab43352</a> Human breast tumour tissue lysate <a href="#">ab43351</a> Human breast normal tissue lysate (matched) Extraction 2: insoluble protein fraction ab43354 Human breast tumour tissue lysate <a href="#">ab43353</a> Human breast normal tissue lysate (matched)</p>

製品の特性

<b>製品の状態</b>	Liquid
<b>保存方法</b>	Shipped on Dry Ice. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.
<b>バッファー</b>	pH: 7.40 Constituents: PBS, 15.22% Thiourea, 30.03% Urea, 0.1% SDS, 0.77% DTT
<b>特記事項 (精製)</b>	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors