

Product datasheet

Anti-CDKN2A/p14ARF antibody ab470

8 References 画像数 3

製品の概要

製品名	Anti-CDKN2A/p14ARF antibody
製品の詳細	Rabbit polyclonal to CDKN2A/p14ARF
由来種	Rabbit
特異性	Specific for p14ARF.
アプリケーション	適用あり: WB, ICC/IF
種交差性	交差種: Human
ポジティブ・コントロール	BT549 cells, HeLa whole cell extract

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	PBS with 0.05% sodium azide
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab470** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/1000. Detects a band of approximately 15 kDa (predicted molecular weight: 14 kDa).
ICC/IF		1/100.

ターゲット情報

関連性

The gene for CDK2NA generates several transcripts/proteins which differ from each other in their first exons. Three of these transcripts are generated by alternative splicing (isoform 1 a.k.a p16INK4A, isoform 2 and isoform 3 a.k.a p12), two of which are known to function as inhibitors of CDK4 kinase. One other transcript that is generated from this gene contains an alternate reading frame (ARF), with the first exon located 20kb upstream of the remainder of the gene(isoform 4 a.k.a. p14ARF, p19ARF, ARF). In spite of the structural and some functional differences, all the proteins encoded by the CDKN2A gene are involved in cell cycle G1 control.


細胞内局在

Cytoplasmic and Nuclear

画像




ab470 diluted 1:267 (7.5 µg/ml) and used on NARF6 cells induced with 1 mM IPTG for 24 hrs to induce p14 ARF.



ab470 used at 1 µg/ml with ECL on human cells transfected with p14ARF construct.

ab470 used at 1 µg/ml with ECL on human cells transfected with p14ARF construct.



All lanes : Anti-CDKN2A/p14ARF antibody (ab470) at 1/400 dilution

Lane 1 : Lysate from ARF negative U2OS cells

Lane 2 : Lysate from ARF positive H1299 cells

Predicted band size: 14 kDa

Observed band size: 15 kDa

The picture shows Western blot using cell lysate from U2OS cells (ARF negative cells, lane 1) and H1299 cells (ARF positive cells, lane 2) stained with ab470 at a dilution of 1/400.

This image was kindly supplied as part of the review submitted by Delin Chen.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors