

Product datasheet

Anti-TIRAP antibody ab47986

画像数 1

製品の概要

製品名	Anti-TIRAP antibody
製品の詳細	Goat polyclonal to TIRAP
特異性	This antibody is expected to recognize both reported isoforms (NP_001034750.1 and NP_683708.1).
アプリケーション	適用あり: ELISA, WB
種交差性	交差種: Human
免疫原	Synthetic peptide: C-QTLLKKPKKRPNSPE , corresponding to internal sequence amino acids 27-41 of Human TIRAP Run BLAST with Run BLAST with
ポジティブ・コントロール	K562 cell lysate.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 0.5% BSA, Tris buffered saline, pH 7.3
精製度	Immunogen affinity purified
特記事項 (精製)	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab47986** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

ELISA

WB

追加情報

Peptide ELISA: Antibody detection limit dilution 1:32,000.

WB: Use at a concentration of 0.1 - 0.3 µg/ml. Detects a band of approximately 24 kDa observed in lysates of Human myeloid erythroblast cell line K562 (predicted molecular weight: 24 kDa).

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報


関連性

The Toll-like receptor (TLR) family in mammals comprises a family of transmembrane proteins characterized by multiple copies of leucine rich repeats in the extracellular domain and IL-1 receptor motif in the cytoplasmic domain. Like its counterparts in Drosophila, TLRs signal through adaptor molecules. The TLR family is a phylogenetically conserved mediator of innate immunity that is essential for microbial recognition. Ten human homologs of TLRs (TLR1-10) have been described. TIRAP (TIR domain-containing adaptor protein) is an adaptor protein used by TLR4. Blocking TIRAP inhibits TLR4-mediated signaling events, including DC maturation and cytokine production. Function: Adapter involved in TLR2 and TLR4 signaling pathways in the innate immune response. Acts via IRAK2 and TRAF-6, leading to the activation of NF-kappa-B, MAPK1, MAPK3 and JNK, and resulting in cytokine secretion and the inflammatory response. Positively regulates the production of TNF-alpha and interleukin-6. Tissue specificity: Highly expressed in liver, kidney, spleen, skeletal muscle and heart. Also detected in peripheral blood leukocytes, lung, placenta, small intestine, thymus, colon and brain. Post-translational modification: Phosphorylated by IRAK1 and IRAK4. Also phosphorylated by BTK. Sequence similarities: Contains 1 TIR domain.

細胞内局在

Cytoplasmic

画像



Anti-TIRAP antibody (ab47986) at 0.1 µg/ml +
K562 cell lysate (35µg protein in RIPA buffer).

Predicted band size : 24 kDa

Observed band size : 24 kDa

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors