

Product datasheet

Anti-EYA2 antibody ab54713

画像数 2

製品の概要

製品名	Anti-EYA2 antibody
製品の詳細	Mouse monoclonal to EYA2
アプリケーション	適用あり: WB, ICC/IF
種交差性	交差種: Human
免疫原	Recombinant fragment, corresponding to amino acids 164-272 of Human EYA2

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None PBS, pH 7.2
精製度	Protein G purified
ポリ/モノ	モノクローナル
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab54713** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 - 5 µg/ml. Predicted molecular weight: 59 kDa.
ICC/IF		Use a concentration of 10 µg/ml.

ターゲット情報

機能

Tyrosine phosphatase that specifically dephosphorylates 'Tyr-142' of histone H2AX (H2AXY142ph). 'Tyr-142' phosphorylation of histone H2AX plays a central role in DNA repair and acts as a mark that distinguishes between apoptotic and repair responses to genotoxic stress. Promotes efficient DNA repair by dephosphorylating H2AX, promoting the recruitment of DNA repair complexes containing MDC1. Its function as histone phosphatase probably explains its role in transcription regulation during organogenesis. Coactivates SIX1. Seems to coactivate SIX2, SIX4 and SIX5. Together with SIX1 and DACH2 seem to be involved in myogenesis. May be involved in development of the eye. Interaction with GNAZ and GNAI2 prevents nuclear translocation and transcriptional activity.

組織特異性

Highest expression in muscle with lower levels in kidney, placenta, pancreas, brain and heart.

配列類似性

Belongs to the HAD-like hydrolase superfamily. EYA family.


発生段階

At the begin of fourth week of development detected in cytoplasm of somite cells. Between the sixth and eighth week of development detected in cytoplasm of limb bud cells.

細胞内局在

Cytoplasm. Nucleus.

画像



Predicted band size : 59 kDa



ab54713 at 10 µg/ml staining EYA2 in human HeLa cells by Immunocytochemistry / Immunofluorescence.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors